



ISSN (E): 2277- 7695  
ISSN (P): 2349-8242  
NAAS Rating: 5.03  
TPI 2019; 8(9): 39-43  
© 2019 TPI  
www.thepharmajournal.com  
Received: 19-07-2019  
Accepted: 23-08-2019

**Surabhi Kumari**  
M.Sc (food tech.) Student,  
Warner College of Dairy  
Technology, SHUATS, Naini,  
Allahabad, Uttar Pradesh,  
India

**Shanker Suwan Singh**  
Assistant Professor, Food  
Technology, WCDT, SHUATS,  
Naini, Allahabad, Uttar,  
Pradesh, India

**Komal Yadav**  
M.Sc (food tech, Student,  
Warner College of Dairy  
Technology, SHUATS, Naini,  
Allahabad, Uttar Pradesh, India

**Bhavya**  
M.Sc food tech, Student,  
Warner College of Dairy  
Technology, SHUATS, Naini,  
Allahabad, Uttar Pradesh,  
India

#### Correspondence

**Surabhi Kumari**  
M.Sc (food tech.) Student,  
Warner College of Dairy  
Technology, SHUATS, Naini,  
Allahabad, Uttar Pradesh,  
India

## Development and quality assessment of Gluten-free Bread prepared by using Rice flour, Corn starch and Sago flour

Surabhi Kumari, Shanker Suwan Singh, Komal Yadav and Bhavya

DOI: <https://doi.org/10.22271/tpi.2019.v8.i9a.3955>

#### Abstract

Baking industry is currently seeking to expand its products range, but also to constitute a way of maintain and improving general health of people. Bakery products are mainly prepared from wheat as its main ingredients. Bread is widely consumed and is an ideal vehicle for functional delivery. The objective of this work was to develop Gluten-free Bread. The use of rice flour, corn starch and sago flour blends as a source of high carbohydrate, protein and fat and absence of gluten in production of Gluten- free bread was studied. The flour blends of rice flour, corn starch and sago flour were composites at replacement levels of 40:35:25 (T<sub>1</sub>), 50:30:20 (T<sub>2</sub>), 60:25:15 (T<sub>3</sub>) % while the wheat flour bread 100:00:00 (T<sub>0</sub>) served as control bread. Various analysis parameters were analyzed by one way ANOVA to obtained a predicted optimum result prepared bread was subjected to chemical, microbial, and sensory analysis to evaluate the suitability of breads were T<sub>1</sub> protein (5.10 %), fat (4.58 %), ash (0.95 %), moisture (31.15 %) and carbohydrate (58.19 %) as comparable to control without adversely affecting the sensory parameters. Based on the result it was indicated that beneficial components of Rice flour, corn starch and sago flour made them more favorable choice for food technologist to develop gluten - free bread especially for celiac disease.

**Keywords:** Rice flour, corn starch, sago flour, celiac disease

#### Introduction

Bread is one of the oldest and largest consumed food stuff and is consumed across the globe by all age groups. Breads are defined as a fermented bakery product produced mainly from wheat flour, water, yeast, and salt by a series of process involving mixing, kneading, fermentation, proofing, shaping and baking (Dewettinck *et al.*, 2008) <sup>[1]</sup>. A gluten-free bread is a bakery product that strictly excludes gluten, a mixture of proteins found in wheat and related grains, including barley, rye, oat, and all their species and hybrids (Such as spelt, kamut, and triticale). Gluten-free breads are made with rice flour, corn starch, sago flour and flours made by other cereals such as sorghum etc. (Biesiekierski, 2017) <sup>[3]</sup>. Gluten causes health problems for those with gluten-related disorders, including coeliac disease (CD), non- coeliac gluten sensitivity (NCGS) and wheat allergy.

Bread products made from wheat flour are consumed worldwide, some individuals are intolerant to prolamins of wheat and the related cereals oats, rye, and barley. This intolerance, celiac disease, seriously impairs intestinal absorption and can lead to severe malnutrition. (Davison *et al.*, 1987; Ciclitira *et al.*, 1987) <sup>[8]</sup>.

The only certain remedy to this pathology is total omission of gluten from the diet. That is very difficult, however, since many foods contain gluten (Baldo *et al.*, 1984) <sup>[4]</sup>. Many investigators have attempted to develop gluten-free yeast breads. Wheat starch has often been used as a wheat flour replacement.

The World Health Organization states that gluten-free foods should contain less than 1 mg gliadin per 100-g product. Non-wheat cereal products may also be used in bread making. Rice flour is useful since it lacks gluten and contains low levels of sodium and a high amount of easily digested carbohydrates, making it desirable in celiac diets.

Celiac disease (CD) is a chronic enteropathy characterized by an inadequate immune response to ingested gluten from wheat, rye, barley, triticale, and, in some cases, oats. This intolerance results in damage to the small-intestinal mucosa, leading to the malabsorption of nutrients. The only effective treatment for CD patients relies on a lifelong gluten-free diet.

The general prevalence of CD is estimated to be one in 300 (Collin *et. al.*, 1997), although recent population-based screening studies suggest that the prevalence may be even higher (one in 100). At previous trials, rice flour, cornstarch, and cassava starch were used separately in preparing bread because these raw materials are normally used by celiac people in food preparation. Nevertheless, better technological results were obtained when these raw materials were used together in the mixture. The aim of the present study was to use rice flour, cornstarch, and cassava starch in gluten-free bread making and to statistically establish optimal amounts of each ingredient.

**Rice flour**

Rice (*Oryza sativa*) flour is a popular and widely used gluten-free flour. Rice flour (Also rice powder) is a form of flour made from finely milled rice. Rice flour may be made from either white rice or brown rice. To make the flour, the husk of rice or paddy is removed and raw rice is obtained, which is then ground to flour. Rice flour contains fat, carbohydrates, protein, calcium, iron. Rice flour is very common in parts of the world where rice is more of a staple ingredient than is wheat or wheat flour. In the 1980's, it was introduced to the western part of the world, where it became a primary ingredient in many processed foods in the baking industry. Of the world's total rice production, 90% is grown and consumed in Asia. White rice flour is enriched with vitamins and minerals to meet the nutrient requirements. Rice flour is gluten-free; therefore, individuals with celiac disease can safely consume it. It is often used as a healthier alternative to wheat flour because it is easier to digest and richer in fiber. Because rice possesses unique nutritional, hypoallergenic, colorless, and bland taste properties, it is used in baby foods and puddings. Its role in development of foods for gluten-intolerant patients especially has been increasing.

**Table 1:** Nutrient Composition of rice Flour

Moisture %	13.3
Protein %	8
Fat %	0.4 - 1
Carbohydrate %	79.1
Fiber %	0.6
Ash %	0.4 - 0.7
Calcium (mg/100g)	10
Phosphorous (mg/100g)	150
Iron (mg/100g)	2.2
Thiamine (mg/100g)	210
Nicotinic acid (mg/100g)	3.8

**Corn starch**

Corn starch or maize starch is the starch derived from the corn (Maize) grain. The starch is obtained from the endosperm of the kernel. Corn starch is a common food ingredient. It is versatile, easily modified, and finds many uses in industry as adhesives, in paper products, as an anti-sticking agent. Corn starch contains carbohydrates, dietary fiber, iron. It has medical uses, such as to supply glucose for people with glycogen storage disease. The corn is steeped for 30 to 48 hours, which ferments it slightly. The germ is separated from the endosperm and those two components are ground separately (still soaked). Next the starch is removed from each by washing. The starch is separated from the corn steep liquor, the cereal germ, the fibers and the corn gluten mostly in hydro cyclones and centrifuges, and then dried. (The

residue from every stage is used in animal feed and to make corn oil or other applications). This process is called wet milling. Finally, the starch may be modified for specific purposes.

**Table 2:** Nutrient composition of corn starch.

Moisture (%)	10
Protein (%)	0.1
Ash (%)	0.2
Fiber (%)	0.1
Fat (%)	0.2
Carbohydrates (%)	87.5

**Sago flour**

Sago (Tapioca Sago) (Sabudana) is a processed, ready to cook agricultural food product. The only raw material for manufacturing Sabudana is "tapioca root" internationally known as "cassava". Sabudana is considered an acceptable form of nutrition. It is used in a variety of dishes such as desserts like "kheer" (Boiled with sweetened milk) or Khichadi, vada, bonda etc. (Mixed with Potatoes, Ground nuts, rock-salt, and black-pepper or green chillies). Sabudana (Sago) is a produce, prepared purely from Tapioca Root ("Cassava") Botanical name is "Mani hot Esculenta Crantz Syn. Utilissima". In India, the names 'Sago', 'Tapioca Sago', 'Tapioca Globules', 'Sabudana', 'Javvarisi', 'Sabbakki', 'Saggubeeyam' are all used for the same commodity 'Sabudana. It provides only carbohydrate food value, and is low in protein, vitamins and minerals. Sago starch can be baked (Resulting in a product analogous to bread, pancake, or biscuit) or mixed with boiling water to form a paste. In many developing countries bread consumption is continually expanding and there is increasing dependence on imported wheat. Most of these countries, however, grow staples other than wheat that can be used for bread. Some grow various starchy tubers such as sago, yam or sweet potatoes and some others grow cereals such as maize, millet or sorghum. It would therefore be economically advantageous for those countries if imports of wheat could be reduced or even eliminated and the demand for bread could be met by the use of domestically grown products instead of wheat. Sago is a common ingredient used in Indian recipes. In gruel form, sago can function as a healthy alternative to carbonated drinks, providing energy without any artificial chemicals and sweetener.

**Table 3:** Nutrient composition of sago

Moisture (%)	13.5
Protein (%)	0.15
Ash (%)	0.73
Fiber (%)	0.4
Fat (%)	0.65
Starch (%)	70.25

**Justification**

The population everywhere desire to eat a healthier diet without changing their conventional dietary patterns. The term gluten-free bread is generally being used for foods that may provide a health and physiological benefits. People decide to go gluten free for a few different reasons – originally, gluten free foods were made available for those suffering from celiac disease, or for those who are sensitive to gluten. More recently a lot of people have decided to go gluten free as they enjoy the taste of well-made items that are free of gluten as

they become more widely available. Celiac disease is a condition in which the mucous membrane of the small intestine of gluten intolerant people is damaged by gluten. The reason gluten can be such an issue for some people is because wheat can be incredibly difficult to digest for some, leaving behind gas in the stomach that can make those with sensitivities feel bloated and uncomfortable. Refined wheat also has very little nutritional value and therefore taking this out of our diet is not usually a problem for us health-wise. Sticking to a gluten free diet can be tricky, and for most of us, cutting bread and cake can be a hard one. Gluten free products allow you to include a little bit of what you enjoy in your diet and also help you feel full throughout the day without the horrible bloated feeling that can come from a sensitivity.

**Review of literature**

Reported that elimination of gluten increases the role and importance of starch in providing structure and texture to gluten-free products. Starch is the primary source of stored energy in many plants including cereals, legumes, roots and tubers. It provides 70%–80% of the calories consumed by humans worldwide. Reported that isolated wheat starch is often utilised in gluten-free products, but starch-based ingredients should ideally originate from raw materials that are naturally gluten-free. Reported that the incorporation of various hydrocolloids into dough formulations has made it possible to produce bread from rice flour. Studied the effect of rice and waxy rice on the French fry-like products using extrusion before frying. Fries from waxy rice flour had texture most comparable to commercial fast food fried potatoes while fries from non-waxy rice flours were too hard and gummy. Reported that similar treatment may also be applied to bakery products although too high content of high amylose starch in bread leads to deterioration of its quality. The similar work for fasting biscuits was conducted by using sago, peanut,


banana, potato, foxtail millet, barnyard millet in different proportions. Reported that Corn ranks as the second most widely produced cereal crop worldwide. Because of the high productivity, corn is by far the most economical cereal to produce. Corn flour contains high levels of many important vitamins and minerals, including potassium, phosphorus, zinc, calcium, iron, thiamine, niacin, vitamin B-6, and folate. Reported that High amylose corn starch is commonly used to increase the level of resistant starch in foods. Gujral *et al.*, (2004) [14] reported that Rice flour exhibits properties such as the absence of gluten, low levels of sodium, protein, fat and fiber, and high amount of easily digested carbohydrates, which are desirable for certain special diets. It has also bland taste, white color, and hypoallergenic properties. Guha *et al.*, (2006) [15] reported that the glutinous rice was suitable material to produce the expanded extruded rice product such as ready-to-eat snacks, breakfast cereal with low bulk density, high expansion and low shear stress. Reported that Rice flour is increasingly popular as a substitute of wheat flour in the preparation of products consumed by wheat-intolerant or celiac patients. It is the most suitable cereal grain flour for the production of gluten-free products due to its bland taste, white colour, digestibility, and hypoallergenic properties.

**Materials and Methods**

**Procurement and collection of ingredients**

- Rice to be purchased from local market of Prayagraj.
- Corn starch to be purchased from local market of Prayagraj.
- Sago to be purchased from local market of Prayagraj.
- Other materials also to be purchased from local market of Prayagraj.

**Flow diagram adopted for preparation of control and experimental bread**



## Result and Discussion

The analyzed data is presented in this chapter under the following headings:

1. Chemical characteristics

2. Microbial characteristics

3. Organoleptic characteristics

4. Statistical characteristics

5. Estimation of cost of production

**Table 4:** Average data for different parameters of control and experiments (in percent)

Parameters	Treatments			
	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>
1. Chemical Analysis				
Moisture	31.72	31.15	31.23	31.36
Fat	4.21	4.58	4.75	4.80
Protein	9.64	5.10	5.60	6.42
Carbohydrate	53.41	58.19	57.50	56.71
Ash	1.00	0.95	0.90	0.68
2. Microbiological analysis				
SPC × 10 <sup>-3</sup> (colony forming unit /gm)	3.8	4.2	4	4.4
Yeast & Moulds count (cfu/gm)	1.4	1.6	1.6	2.2
Coliform count	Nil	Nil	Nil	Nil
3. Organoleptic Score (9- point hedonic scale)				
Color and Appearance	7.54	8.08	7.39	6.7
Body and Texture	7.85	8.04	7.44	6.76
Flavor and taste	7.73	8.16	7.4	7.12
Overall acceptability	7.70	8.09	7.40	6.85
4. Cost analysis				
Cost in Rs./100g	5.87	41.25	36.36	31.45

## Summary and Conclusion

### Physico- chemical analysis

#### Moisture percentage

There was significant difference in moisture content of different treatment combinations. Maximum moisture percent was recorded in the sample of T<sub>0</sub> (31.76) followed by T<sub>3</sub> (31.36), T<sub>2</sub> (31.24), T<sub>1</sub> (31.15). The difference in moisture was due to the composition difference of wheat flour, rice flour, corn starch and sago flour which are used in different proportions in different treatments.

#### Ash percentage

There was significant difference in ash content of different treatment combinations. Maximum ash percent was recorded in the sample of T<sub>0</sub> (1.01) followed by T<sub>1</sub> (0.96), T<sub>2</sub> (0.90), T<sub>3</sub> (0.68). The difference in ash was due to the composition difference of wheat flour, rice flour, corn starch and sago flour which are used in different proportions in different treatments.

#### Protein percentage

There was significant difference in protein content of different treatment combinations. Maximum protein percent was recorded in the sample of T<sub>0</sub> (9.64) followed by T<sub>3</sub> (6.43), T<sub>2</sub> (5.61), T<sub>1</sub> (5.11). The difference in protein was due to the composition difference of wheat flour, rice flour, corn starch and sago flour which are used in different proportions in different treatments.

#### Fat percentage

There was significant difference in fat content of different treatment combinations. Maximum fat percent was recorded in the sample T<sub>3</sub> (4.81) followed by T<sub>2</sub> (4.75), T<sub>1</sub> (4.59), T<sub>0</sub> (4.21). The difference in fat was due to the composition difference of wheat flour, rice flour, corn starch and sago flour which are used in different proportions in different treatments.

#### Carbohydrate percentage

There was significant difference in carbohydrate content of

different treatment combinations. Maximum carbohydrate percent was recorded in the sample of T<sub>0</sub> (31.76) followed by T<sub>1</sub> (58.20) followed by T<sub>2</sub> (57.50), T<sub>3</sub> (56.71), T<sub>0</sub> (53.42). The difference in carbohydrate was due to the composition difference of wheat flour, rice flour, corn starch and sago flour which are used in different proportions in different treatments.

### Organoleptic analysis

#### Color and appearance score

There was significant difference in color and appearance score of different treatment combinations. Maximum color and appearance score was recorded in the sample of T<sub>0</sub> (31.76) followed by T<sub>1</sub> (8.08) followed by T<sub>0</sub> (7.54), T<sub>2</sub> (7.39), T<sub>3</sub> (6.70). The difference in colour and appearance was due to the composition difference of wheat flour, rice flour, corn starch and sago flour which are used in different proportions in different treatments.

#### Body and texture

There was significant difference in body and texture score of different treatment combinations. Maximum body and texture score was recorded in the sample of T<sub>0</sub> (31.76) followed by T<sub>1</sub> (8.04) followed by T<sub>0</sub> (7.84), T<sub>2</sub> (7.44), T<sub>3</sub> (6.76). The difference in body and texture was due to the composition difference of wheat flour, rice flour, corn starch and sago flour which are used in different proportions in different treatments.

#### Flavor and taste

There was significant difference in flavor and taste score of different treatment combinations. Maximum flavor and taste score was recorded in the sample of T<sub>0</sub> (31.76) followed by T<sub>1</sub> (8.16) followed by T<sub>0</sub> (7.73), T<sub>2</sub> (7.54), T<sub>3</sub> (7.24). The difference in flavor and taste was due to the composition difference of wheat flour, rice flour, corn starch and sago flour which are used in different proportions in different treatments.

### Overall acceptability

There was significant difference in overall acceptability score of different treatment combinations. Maximum overall acceptability was recorded in the sample of T<sub>0</sub> (31.76) followed by T<sub>1</sub> (8.09) followed by T<sub>0</sub> (7.69), T<sub>2</sub> (7.41), T<sub>3</sub> (6.85). The difference in overall acceptability was due to the composition difference of wheat flour, rice flour, corn starch and sago flour which are used in different proportions in different treatments.

### Microbiological parameters

#### Yeast and mould count (cfu/gm)

The highest mean value for yeast and mould count of bread for control and treatment samples was recorded and was significantly viable as T<sub>3</sub> (2.20) followed by T<sub>1</sub> (1.60), T<sub>2</sub> (1.60) and the lowest is T<sub>0</sub> (1.40).

#### Coliform count

It is evident from the experiment that the coliform test of control and experimental sample was 100 percent negative.

#### Standard plate count

The highest mean value for standard plate count of bread for control and treatment samples was recorded and was significantly viable as T<sub>3</sub> (4.40) followed by T<sub>1</sub> (4.20), T<sub>2</sub> (4.00) and the lowest is T<sub>0</sub> (3.80).

### Conclusion

The present experiment entitled "Development and Quality Assessment of Gluten-Free Bread Prepared by Using Rice flour, Corn Starch and Sago flour" was conducted for the preparation of bread using rice flour, corn starch and sago flour in the proportions, (40:35:25), (50:30:20), (60:25:15). Thus final product was analyzed for their Physico-chemical, Microbiological, sensory and cost. Our results show, therefore, that high-quality, gluten-free bread with good taste and appearance, suitable for celiac patients, can be made from a mixture of 3 raw materials normally consumed in other foods: cornstarch, rice flour, and sago flour. The indicative optimal formulation for this bread was (T<sub>1</sub>) rice flour 40%, corn starch 35% and sago flour 25% and its validation will be made with a larger panel.

### References

1. Alpaslan M, Hayta M, The effects of flaxseed, soy and corn flours on the textural and sensory properties of a baked product. *Journal of Food Quality*. 2006; 29:617-627.
2. Abdel-Aal E.-SM. Functionality of Starches and Hydrocolloids in Gluten-Free Foods. In *Gluten-Free Food Science and Technology*; Gallagher E, Ed; Wiley-Blackwell: Oxford, UK, 2009, 200.
3. Biesiekierski JR "What is gluten?" *J. Gastroenterol Hepatol (Review)*. 32 Suppl. 2017; 1:78-81.
4. Baldo BA, Wrigley CW. Allergies to cereals. In: *Pomeranz Y. Advances in cereal science and technology*. St. Paul, Minn AACC. 1984; 6:331-344.
5. Bryant RJ, Kadan RS, Champagne TE, Vinyard BT, Boykin D. Functional and digestive characteristics of extruded rice flour. *Cereal Chemistry*. 2001; 78:131-137.
6. Bean MM, Nishita KD. Rice flours for baking. In *Juliano BO, editor. Rice: chemistry and technology*, 2nd ed. St. Paul: Amer Assoc of Cereal Chemists, 1985, 539-556.
7. Chrastil J. Correlation between the physicochemical and

- functional properties of rice. *J Agric Food Chem*. 1992; 40(6):1683-1686.
8. Ciclitira PJ, Ellis HJ. Investigation of cereal toxicity in coeliac disease. *Postgrad Med J*. 1987; 63:767-775.
9. Davison AGF, Bridges M. Coeliac disease. *Clin Chim Acta*. 1987; 163:1-40.
10. Dewettinck K, Van Bockstaele F, Kuhne B, Vande Walle, Courtens T, Gellynck X. Nutritional value of bread: Influenced of processing, food interaction and consumer perception. *Review Journal Cereal Science*. 2008; 48:243-257.
11. Dervas G, Doxastakis G, Hadjisavva-Zinoviadi S, Triantafyllakos N. Lupin flour addition to wheat flour dough and effect on rheological properties. *Food Chemistry*. 1999; 66:67-73.
12. Dhingra S, Jood S. Organoleptic and nutritional evaluation of wheat breads supplemented with soybean and barley flour. *Food Chemistry*. 2001; 77:479-488.
13. Eliasson A, Larsson K, *Cereals in Breadmaking: A Molecular Colloidal Approach*. New York, NY: Marcel Dekker, 1993.
14. Gujral S, Haros M, Rosell C. Improving the texture and delaying staling in rice flour chapatti with hydrocolloids and  $\alpha$ -amylase. *Journal of Food Engineering*. 2004; 65:89-94.
15. Guha M, Ali SZ. Extrusion cooking of rice effect of amylose content and barrel temperature on product profile. *Journal of food processing and preservation*. 2006; 30:706-716.
16. Hung PV, Yamamori M, Morita N. Formation of enzyme-resistant starch in bread as affected by high-amylose wheat flour substitutions. *Cereal Chemistry*. 2005; 82:690-694.
17. Moore MM, Schober TJ, Dockery P, Arendt EK. Textural comparisons of gluten-free and wheat-based doughs, batters, and breads. *Cereal Chemistry*. 2004; 81(5):567-575.
18. Miyazaki MR, Hung PV, Maeda T, Morita N. Recent advances in application of modified starches for breadmaking. *Trends Food Sci. Technol*. 2006; 17:591-599.
19. Nishita KD, Bean MM. Physico-chemical properties of rice in relation to rice bread. *Cereal Chemistry*. 1979; 56:185-189.
20. Navickis L. Corn flour addition to wheat flour dough-effect on rheological properties. *Cereal Chemistry*. 1987; 64:307-310.