



ISSN (E): 2277- 7695
ISSN (P): 2349-8242
NAAS Rating: 5.03
TPI 2018; 7(4): 91-93
© 2018 TPI
www.thepharmajournal.com
Received: 08-02-2018
Accepted: 09-03-2018

Rekha Devi
Department of Livestock
Products Technology, College of
Veterinary Science, Lala Lajpat
Rai University of Veterinary and
Animal Sciences, Hisar,
Haryana, India

Nita Khanna
Department of Livestock
Products Technology, College of
Veterinary Science, Lala Lajpat
Rai University of Veterinary and
Animal Sciences, Hisar,
Haryana, India

Vaquil
Department of Livestock
Products Technology, College of
Veterinary Science, Lala Lajpat
Rai University of Veterinary and
Animal Sciences, Hisar,
Haryana, India

Apoorva Argade
Department of Livestock
Products Technology, College of
Veterinary Science, Lala Lajpat
Rai University of Veterinary and
Animal Sciences, Hisar,
Haryana, India

SS Ahlawat
Department of Livestock
Products Technology, College of
Veterinary Science, Lala Lajpat
Rai University of Veterinary and
Animal Sciences, Hisar,
Haryana, India

Correspondence
SS Ahlawat
Department of Livestock
Products Technology, College of
Veterinary Science, Lala Lajpat
Rai University of Veterinary and
Animal Sciences, Hisar,
Haryana, India

Utilization of soy milk in strawberry pulp based shrikhand for development of a novel fermented milk product

Rekha Devi, Nita Khanna, Vaquil, Apoorva Argade and SS Ahlawat

Abstract

The present investigation was undertaken to explore the possibility of the use of soy milk in strawberry pulp based shrikhand to produce a novel fermented milk product. The product was prepared as per the standard method using constant level of strawberry pulp (15%) and sugar (40%). Different levels of soy milk (10, 20, 30 and 40%) were tried as substitute of standardized milk. It was found that incorporation of soy milk at 20% level slightly improved the colour and appearance scores of the developed product. Other sensory parameter (Flavour, Body and texture, Sweetness and Overall Acceptability) scores were decreased as the level of soy milk incorporation increased, but the scores were comparable with control samples at 20% soy milk incorporation. The moisture and protein content was increased and fat, titrable acidity and total solids were decreased in 20% soy milk incorporated developed shrikhand. It is concluded that strawberry pulp (15%) based shrikhand can be prepared by replacing 20% soy milk with standardized milk without compromising its quality.

Keywords: shrikhand, strawberry pulp, standardized milk, soy milk, quality

Introduction

Fermented milk products constitute a vital component of the human diet in many regions of the world [1]. Indian fermented milk products utilize 7 per cent of total milk produced and mainly include three sweetened products i.e. dahi, shrikhand and lassi. These products have enjoyed reputation for their nutritional and therapeutic value from time immemorial [2] and play an important role in synthesis of vitamin B complex in human body. These products also prevent the stomachic diseases, because several lactic organisms produce natural antibiotics [3]. Shrikhand is popular dessert and forms part of a delicious supplement on religious functions, particularly in the state of Maharashtra, Gujarat, Karnataka and some parts of South India [2].

With the restricted availability and high cost of animal milk, the scientists have been making attempts to switch over to the utilization of plant proteins. Soya bean is nutritious and used to prepare soy milk based fermented food products. Fermentation of soy milk improves the flavour, enhances the nutritional quality of food, increases the digestibility, eliminates the anti nutritional factors, prolongs shelf life, adds therapeutic value and is relatively economic as compared to dairy milk products [4].

Soy milk based fruit juice beverage would offer several distinct nutritional advantages over the plain fruit beverage to the consumer. It is also a common practice of using fruits in preparation of various dairy products like ice cream, yoghurt and shrikhand [3]. Fruit pulp provides sweetness and masks the beany flavor of soy milk to some extent [5]. Various workers have tried pulp of various fruits in the preparation of shrikhand [6].

Hence, the present investigation was undertaken to explore the possibility of the use of soy milk in strawberry pulp based shrikhand to produce a novel fermented milk product.

Materials and Methods

Soy bean, strawberry fruits and powdered Sugar were purchased from local market and fruit pulp was used for enhancing the flavour and acceptability of product. Standardized milk having 4.5% fat and 8.5% SNF was procured from experimental dairy plant, department of Livestock Products Technology, COVS, LUVAS, Hisar and the mixed starter culture NCDC-263 (*Streptococcus thermophilus* and *Lactobacillus bulgaricus*) was procured from National Collection of Dairy Cultures, NDRI, Karnal.

The strawberry pulps were prepared with standard methods in hygienic conditions and

pasteurization was done at 76 °C for 1 min. The pulp were packaged in polythene bags and sealed aseptically. Soy milk was prepared in the College of Agriculture Engineering, CCS, HAU Hisar as per standard procedures [7] and packed in glass bottles. Both the ingredients were shifted to cold storage (4±1 °C) room for further use [8].

Treatments were prepared by replacing standardized milk with soy milk at different levels (10, 20, 30 and 40%). Strawberry pulp (15%) and sugar (40%) was added to both control and treated samples as per the standardized method [3, 9] and further processed for preparation of shrikhand as per the method followed by Sonawane [3].

Treatments

- Control (C): Chakka prepared from 100% standardized milk.
- T₁: Chakka prepared from 90% standardized milk + 10% soy milk.
- T₂: Chakka prepared from 80% standardized milk + 20% soy milk.
- T₃: Chakka prepared from 70% standardized milk + 30% soy milk.
- T₄: Chakka prepared from 60% standardized milk + 40% soy milk.

Method of Trout [10] was followed for determining the pH of the samples and the acidity (titration) and proximate composition of sample were determined by following the method described by AOAC [11].

The nine point hedonic scale was employed for sensory evaluation following the methodology described by Harry and Hildegard [12].

The experiment was replicated thrice along with duplicate samples and data obtained were subjected to one way ANOVA followed by Duncans Multiple Range Test by using SPSS software [13].

Results and Discussion

Sensory Evaluation

The colour and appearance score was slightly improved with addition of soy milk at T₁ and T₂ treatment levels as compared to control (C) for development of strawberry based shrikhand (Table 1), however, it was significantly ($p \leq 0.05$) decreased with further incorporation of soy milk (T₃ and T₄) as compared to control as well as T₁ and T₂ samples. The colour and appearance scores of T₂ were statistically comparable with C, and hence selected. These results of colour and appearance for soya milk incorporation are in accordance with the findings of Deshpande [7, 14, 15], who reported lower average colour and appearance in shrikhand prepared from buffalo milk blended at higher levels of soy milk as compared to control.

It was evident from the results that the flavor score was decreased as the level of soy milk increased. It was due to beany flavor of soy milk as reported by Deshpande *et al.* [16]. The treatment T₂ was selected by judges because it contain higher proportion of soy milk than T₁ with acceptable scores and statistically comparable with control samples. The lower flavor score in soy milk incorporated shrikhand are in agreement with the results of Sumedha *et al.* [15] and Borate *et al.* [17], who reported lower average flavor score in shrikhand prepared from buffalo milk blended at higher levels of soy milk due to its dominating beany flavor as compared to control.

Table 1: Effect of soy milk incorporation on sensory score of strawberry pulp based shrikhand.

Sensory attributes	C	T ₁	T ₂	T ₃	T ₄
Colour and appearance	7.83 ^a ±0.16	8.00 ^a ±0.00	8.00 ^a ±0.00	7.23 ^b ±0.21	7.00 ^b ±0.36
Flavour	8.08 ^a ±0.25	7.83 ^a ±0.16	7.75 ^a ±0.20	7.00 ^b ±0.12	6.42 ^c ±0.20
Body and texture	8.33 ^a ±0.32	7.92 ^a ±0.83	8.08 ^a ±0.20	7.08 ^b ±0.08	6.42 ^c ±0.20
Sweetness	8.00 ^a ±0.25	7.83 ^a ±0.16	7.83 ^a ±0.16	7.16 ^b ±0.10	6.33 ^c ±0.21
Overall acceptability	7.83 ^a ±0.16	7.73 ^a ±0.10	7.75 ^a ±0.17	6.67 ^b ±0.16	6.00 ^c ±0.00

Mean±SE, with different superscript in a row differ significantly ($p \leq 0.05$) (n=18)

C=Control, T₁: 90%STD milk+10%soy milk, T₂: 80%STD milk+20%soy milk.

T₃: 70%STD milk+30%soy milk,

T₄: 60%STD milk+40%soy milk.

The body and texture score was slightly decreased with addition of soy milk at T₁ and T₂ treatment levels as compared to control (C) for development of strawberry based shrikhand, however, it was significantly ($p \leq 0.05$) decreased with further incorporation of soy milk as compared to control as well as T₁ and T₂ samples. The body and texture scores of T₂ were statistically comparable with C, and hence selected. Deshpande *et al.* [16] also reported the similar results for body and texture scores with incorporation of soy milk in standardized buffalo milks shrikhand. The lower body and texture scores in soy milk incorporated shrikhand are also in agreement with the results of Sumedha *et al.* [15] and Barate *et al.* [17].

However, the sweetness scores were decreased as the level of soy milk incorporation increased, but the significant decline was recorded at T₃ and T₄ levels of treatments as compared to C including T₁ and T₂ treatments. The sweetness score of treatment T₂ was comparable with control and hence selected. These results of soy milk incorporated shrikhand might be due to higher acidity and lower average sweetness scores in

shrikhand prepared from buffalo milk blended with soy milk as compared to control samples [15, 17].

The overall acceptability score was slightly decreased with addition of soy milk at T₁ and T₂ treatment levels as compared to control for development of strawberry based shrikhand, however, it was significantly ($p \leq 0.05$) decreased with further incorporation of soy milk at T₃ and T₄ as compared to C as well as T₁ and T₂ samples. The overall acceptability scores of T₂ treatment were statistically comparable with C. The lower acceptability scores at higher level of soy milk incorporated shrikhand are also in agreement with the results of Sumedha *et al.* [15] and Barate *et al.* [17].

On the basis of sensory evaluation, the strawberry (15%) based shrikhand incorporated with 20% soya milk was selected for further studies

Physico-chemical properties

Moisture content strawberry (15%) based shrikhand was significantly ($P \leq 0.05$) increased with addition of 20% soy milk (Table 2). Total solids of control was significantly

($P \leq 0.05$) higher than soya milk added product. This might be due to the fact that soy proteins have high affinity for water binding and thus having high water holding capacity. Similar

results were also reported by Deshpande *et al.* [7, 14] and Borate *et al.* [17].

Table 2: Physico Chemical properties of soy milk blended strawberry pulp based shrikhand.

Parameters	Treatment	
	Strawberry based Shrikhand (Control)	Strawberry based Shrikhand with 20% Soya milk
Moisture (%)	51.73 ^b ± 0.24	54.87 ^a ± 0.11
Total Solids (%)	48.27 ^a ± 0.21	45.13 ^b ± 0.32
Protein (%)	5.60 ^b ± 0.03	6.47 ^a ± 0.04
Fat (%)	8.92 ^a ± 0.01	7.12 ^b ± 0.08
Ash (%)	0.57 ± 0.03	0.55 ± 0.02
Sucrose (%)	24.92 ± 0.72	24.29 ± 0.83
pH	3.71 ± 0.02	3.74 ± 0.05
Titration acidity (%)	1.62 ^a ± 0.01	1.45 ^b ± 0.04

Mean ± SE with different superscript in a row differ significantly ($p \leq 0.05$). (n=6)

There was a significant ($P \leq 0.05$) increase in protein content of soya milk added shrikhand as compared to control. Present findings for protein content of soy milk blended shrikhand are in line with that reported by Borate *et al.* [17]. Fat content of soya milk added shrikhand was significantly ($P \leq 0.05$) lower than control. It might be due to more water retained in soy milk added shrikhand which affected the proportionate nutritional parameters [7, 14, 17].

Ash, sucrose content and pH level did not show any significant ($P \leq 0.05$) differences between control and soya milk treated shrikhand and the pH value of developed shrikhand and control sample were within the range prescribed by Datta and Ghatak [18].

The per cent titration acidity of control shrikhand was significantly ($P \leq 0.05$) higher than that of titration acidity of soya milk added developed product. Lower acidity in the fresh shrikhand prepared from buffalo milk blended with soy milk has been reported than shrikhand prepared only from buffalo milk [17]. However, the titration acidity in both control and soya treated products was found well within the permissible limits of 1.4 per cent as per the BIS specification for shrikhand.

Conclusion

It is concluded that strawberry pulp based shrikhand can be prepared with incorporation of 20% soy milk without compromising the sensory, nutritional and physico-chemical quality of the developed shrikhand.

References

1. Swapna G, Chavannavar SV. Shrikhand – Value added traditional dairy product. *Inter. J Food Nutri. Sci.* 2013; 2(4):45-51.
2. Aneja RP, Mathur BN, Chandan RC, Banerjee AK. *Technology of Indian milk products.* Dairy India Publication, Delhi, India, 2002.
3. Sonawane VM, Chavan KD, Pawar BK. Effect of levels of strawberry pulp and sugar on chemical composition during storage of shrikhand. *J Dairying Foods and Home Sci.* 2007; 26(3, 4):153-158.
4. Hasseltine CW, Wanghua HL. The importance of traditional fermented foods. *Bioscience.* 1980; 30:402-404.
5. Lee SY, Morr CV, Seo A. Comparison of milk based and soymilk based yogurt. *J Food Sci.* 1990; 55:532-536.
6. Gavane PF, Zinjarde RM, Rokde SN. Studies on preparation of shrikhand blended with custard apple pulp-

A new fermented milk product. *Indian J Dairy Sci.* 2010; 63(1):11-15.

7. Deshpande S, Bargale PC, Jha K. Suitability of soy milk for development of shrikhand. *J Food Sci. Technol.* 2008; 45(3):284-286.
8. Gandhi NM, Jain SC. A study on the development of a new high protein formulated food using buffalo milk. *J. Food Sci. Technol.* 1977; 14:156.
9. Thakur SN, RajniKant, Chandra R. Preparation of Shrikhand by using mango pulp. *Bioved.* 2014; 25(1):79-82.
10. Trout ES, Hunt MC, Jhonson DE, Clans JR, Castner CL, Kropt DH. Characteristics of low fat ground beef containing texture modifying ingredients. *J Food Sci.* 1992; 57:19.
11. AOAC. *Official Methods of Analysis*, 17th edition, Association of Official Analytical Chemists, Washington, DC, 2002.
12. Harry TL, Hildegard H. *Sensory evaluation of food.* Chapman and Hall, New York, 1998.
13. Snedecor GW, Cochran WG. *Statistical Methods.* VIII edn. Oxford and IBH Publishing Co., New Delhi, 1994.
14. Deshpande S, Bargale PC, Jha K. Quality and acceptability of soy milk based amrakhand. *J. Dairying, Foods and Home Sci.* 2005; 24(1):16-19.
15. Deshpande S, Bargale PC, Jha K. Suitability of soy milk for development of shrikhand. *J Food Sci. Technol.* 2008; 45(3):284-286.
16. Deshpande S, Bargale PC, Jha K. Soy amrakhand – Nutritive food product. *Udyamita.* 2006; 10:46.
17. Borate AJ, Gubbawar SG, Shelke RR, Chavan SD. Studies on keeping quality of Shrikhand prepared from buffalo milk blended with soy milk. *Food Sci. Res. J.* 2011; 2(2):205-210.
18. Datta S, Ghatak PK. Shrikhand-A review. *J Dairy Food and Home Sci.* 2001; 20:141-148.