



ISSN (E): 2277- 7695
ISSN (P): 2349-8242
NAAS Rating: 5.03
TPI 2019; 8(9): 168-170
© 2019 TPI
www.thepharmajournal.com
Received: 07-07-2019
Accepted: 09-08-2019

Shveta Bathla
Deenbandhu Chhotu Ram
University of Science and
Technology, Sonapat, Haryana,
India

Anil Sindhu
Deenbandhu Chhotu Ram
University of Science and
Technology, Sonapat, Haryana,
India

Shivam Kr. Dubey
National Dairy Research
Institute, Karnal, Haryana,
India

Sudarshan Kumar
National Dairy Research
Institute, Karnal, Haryana,
India

Ashok Kumar Mohanty
National Dairy Research
Institute, Karnal, Haryana,
India

Correspondence
Sudarshan Kumar
National Dairy Research
Institute, Karnal, Haryana,
India

Comparative identification of pathogenic bacteria in an infected Karan Fries cow milk

Shveta Bathla, Anil Sindhu, Shivam Kr. Dubey, Sudarshan Kumar and Ashok Kumar Mohanty

Abstract

Mastitis is a multi-etiological disease of the mammary gland characterized mainly by reduction in milk production and milk quality due to intramammary infection by pathogenic bacteria. Nearly 83% of lactating dairy cows in India are infected with mastitis with various degree of inflammation. In this study, we determined differential count of somatic cell count, bacterial load and presence of *S. aureus* in milk collected from healthy, sub-clinical (SCM) and clinical mastitis (CM) Karan Fries cows. On the basis of SCC and CMT score milk was categorized into Healthy, SCM and CM. The SCM and CM milk samples were positive for *S. aureus* on MSA and Baird Parker Agar media while healthy milk samples were negative for *S. aureus*.

Keywords: Intramammary infection, sub-clinical mastitis, clinical mastitis, somatic cell count, california mastitis test, bacteriological

Introduction

Milk is largely made up of water, within which a wide range of nutrients including vitamins, proteins, fats and carbohydrates are suspended [1]. These rich nutritional contents and the production and processing procedures in commercial milk production render it susceptible to contamination by a host of pathogenic microbes that could cause diseases in humans. Therefore, milk is known to be an efficient vehicle for transmission of disease-causing agents to humans [2] and represents the serious health threat to consumers worldwide. The microbial growth has negative impact on product productivity and quality [3].

Mastitis remains one of the most prevalent reason of bacterial load in milk and economically detrimental diseases in dairy farms worldwide [4].

The inflammation of the mammary gland predominantly caused by coagulase-negative *Staphylococci*, *Bacillus* spp., *Streptococcus* spp., *Staphylococcus aureus* (*S. aureus*), and *Escherichia coli* (*E. coli*) [5]. Therefore, the purpose of present study to identify the pathogenic bacteria in milk collected from healthy, sub-clinical infected and clinical infected milk. The bacteriological identification in milk allows to detect the causative agent of disease in herd and helps to provide preventive treatment at accurate time.

Material and Method

Milk collection

Milk sample were collected on quarter basis from 25 Karan Fries (quarter-100) cows from Cattle yard of ICAR-National Dairy Research Institute. The animal selected for milk collection were from various lactation period namely such as early, mid and late stages (50-250).

Determination of Somatic cell count

The somatic cell count was determined by using Eko-milk Scan (Eon Trading LLC, USA) within two hour of milk collection. The milk (5ml) and detergent solution (10ml) was poured in flask mounted on sample mixture in 1:2 ratio and allowed to mix thoroughly by rotational mixing using 10 cycles of swings. After mixing sample was dripped off as a gel and somatic count were displayed on screen.

California Mastitis test

California Mastitis Test (CMT) was performed as prescribed by Schalm and Noorlander (1957) with minor modifications [6]. 1ml of milk from each quarter was poured into four well

paddle. Equal amount (1ml) of CMT reagent was added into each well. The gentle shaking was done for proper mixing and observe the gel/flakes formation in each well. The wells were scored 0 to 3 on the basis of gel formation.

Determination of Bacterial Count

The Standard Plate count method was used for detection of bacterial load in milk collected from healthy, subclinical and clinical mastitis animals. 1ml of milk from each category was added into 5ml of LB Broth at 37 °C for overnight enrichment.

Then enriched bacteria in milk samples were diluted (10⁻¹ to 10⁻⁹) using peptone water and plated on LB media and incubate at 37 °C for 24 hrs.

Detection of Staphylococcus Aureus in milk

Mannitol Salt Agar for the detection of Staphylococcus aureus

A loopful enriched milk from each group was plated on solidified MSA media and incubated at 37 °C for 48 hrs. The plates were observed for media color change and identify the presence of *S. aureus*.

Detection of Staphylococcus aureus on Baird Parker agar

1ml of enriched milk from each group was plated on solidified media and incubated at 37 °C for 48 hrs. The plates were observed for black/grey color selective colonies of *S. aureus*.

Results and Discussion

The SCC is considered as important breast health index in cows and primary indicator of infection. The significant increase of SCC in milk associated poor quality of milk due to low level of lactose, protein, fat and relatively high level of milk pH, sodium and chloride content (electrical conductivity).

The milk samples were categorized by SCC threshold for this study was: Healthy (7x10⁵- 1x10⁶ cells/ml), SCM (2-3.5x10⁶ cells/ml) and CM (13-15x10⁶ cells/ml).

Out of 100 samples, 35% samples had no gel formation and scored 0 indicates the non-infective samples. In 24% samples minor gel formation observed and scored as 1&2 showing sub-acute infection. But in 37% samples solid gel formation was observed and scored as 3&4 which confirmed the severe infection in udder. The correlation of SCC and CMT shown in Table 1.

Table 1: The correlation of SCC and CMT

Health Status	SCC	CMT
Healthy	70-1 lac	0
SCM	2-5 lac	+++
CM	13-15 lac	+++

The enhanced somatic cell count was observed in lipopolysaccharide and lipoteichoic-acid-induced mastitis as compare to healthy bovine milk [7]. The effect of increased SCC on milk composition such protein, lactose, fat and SNF was evaluated by Reis *et al.* (2013). This study showed that IMI has negatively effects on milk composition, although it has been shown that the degree of changes depends on the inflammatory response, the severity and amount of affected tissue in the mammary gland, and bacterial pathogenicity [8]. The healthy, SCM and CM milk were showing differential bacterial load. The bacterial count in healthy milk was less

than 30 cfu/ml and for subclinical and clinical milk it was 30-300 cfu/ml and > 300 cfu/ml respectively shown in Fig.




Fig 1: Detection of Bacterial Load Milk Collected from (A) Healthy (B) Sub-clinical Mastitis (SCM) (C) Clinical Mastitis (CM)

The presence of mastitis causative agent *S.aureus* was confirmed using Mannitol Salt Agar and Baird Parker agar media. In both selective media, healthy samples was showing no growth of *S.aureus* but SCM and CM were showing *S.aureus* growth on MSA and BPA media shown in fig. a and b




Fig 2 (A, B): Selective Detection of *S.aureus* in Healthy, Sub-clinical Mastitis (SCM) and Clinical Mastitis (CM) milk on (A) Mannitol Salt Agar (B) Baird Parker agar media

Harjanti *et al.* (2018) reported that the bacteriological examination of milk samples revealed that Streptococcus was predominant species (73.3%) and the coagulase negative Staphylococcus species was identified at the least bacteria (26.7%) in analyzed samples [9].

Another study reported that 87 staphylococcal isolates were found in milk collected from sub clinical mastitis buffalo, bovine ovine and caprine. Total 58 isolates were positive for mannitol fermentation and remaining 30 isolates produced agglutination with the Staphytech plus kit leads to acetoin production and were classified as *S. aureus* [10].

Furthermore, methicillin-resistant *Staphylococcus aureus* (MRSA) was found in bovine mastitis. The *mecA* gene present in MRSA bacteria confers resistance to almost all β -lactam antibiotics which are the most frequent drugs used in bovine mastitis therapy [11].

Conclusion

In this present study, IMI resulted in increased SCC which was associated with reduced milk production and poor quality of milk. The enhanced SCC reflects the elicited immune response due to invasion of pathogen. More studies are necessary to better understand the variation of mastitis pathogens according to the age, parity, lactation stage and infection stage of the cow.

References

1. Donkor ES, Aning KG, Quay J. Bacteriological Contamination of Informally Marketed Milk in Ghana. *J Food Sci.* 2007; 41:159-174.
2. Donkor ES, Aning KG, Quay J. Bacteriological Contamination of Informally Marketed Milk in Ghana. *J Food Sci.* 2007; 41:159-174.
3. Tauxe Robert V *et al.* Evolving public health approaches to the global challenge of foodborne infections. *International journal of food microbiology.* 2010; 139:S16-S28.
4. Hagnestam-Nielsen, Christel, Søren Østergaard. Economic impact of clinical mastitis in a dairy herd assessed by stochastic simulation using different methods to model yield losses. *Animal.* 2009; 3.2:315-328.
5. Levison LJ *et al.* Incidence rate of pathogen-specific clinical mastitis on conventional and organic Canadian dairy farms. *Journal of dairy science.* 2016; 99.2:1341-1350.
6. Schalm OW. Experiments and observations leading to development of the California Mastitis Test. *J. Am. Vet. Med. Assoc.* 1957; 130:199-204.
7. Wall Samantha K *et al.* Differential somatic cell count in milk before, during, and after lipopolysaccharide-and lipoteichoic-acid-induced mastitis in dairy cows. *Journal of dairy science.* 2018; 101.6:5362-5373.
8. Dos Reis, Carolina Barbosa Malek *et al.* Effect of somatic cell count and mastitis pathogens on milk composition in Gyr cows. *BMC veterinary research.* 2013; 9.1:67.
9. Harjanti DW *et al.* Isolation and identification of bacterial pathogen from mastitis milk in Central Java Indonesia. *IOP Conference Series: Earth and Environmental Science.* IOP Publishing, 2018; 102(1).
10. Aires-de-Sousa, Marta *et al.* Characterization of *Staphylococcus aureus* isolates from buffalo, bovine, ovine, and caprine milk samples collected in Rio de Janeiro State, Brazil. *Appl. Environ. Microbiol.* 2007; 73.12:3845-3849.
11. Srednik Mariela E *et al.* First isolation of a methicillin-resistant *Staphylococcus aureus* from bovine mastitis in Argentina. *Veterinary and Animal Science.* 2019; 7:100043.