



ISSN (E): 2277- 7695

ISSN (P): 2349-8242

NAAS Rating: 5.03

TPI 2019; 8(9): 316-323

© 2019 TPI

www.thepharmajournal.com

Received: 25-07-2019

Accepted: 27-08-2019

K Harish

Department of Veterinary Public Health & Epidemiology
College of Veterinary Science,
Tirupati, Andhra Pradesh, India

A Jagadeesh Babu

Department of Veterinary Public Health & Epidemiology
College of Veterinary Science,
Tirupati, Andhra Pradesh, India

T Madhava Rao

Department of Veterinary Public Health & Epidemiology
College of Veterinary Science,
Tirupati, Andhra Pradesh, India

B Somasekhara Goud

Department of Veterinary Public Health & Epidemiology
College of Veterinary Science,
Tirupati, Andhra Pradesh, India

Y Suresh

Department of Veterinary Public Health & Epidemiology
College of Veterinary Science,
Tirupati, Andhra Pradesh, India

Correspondence

K Harish

Department of Veterinary Public Health & Epidemiology
College of Veterinary Science,
Tirupati, Andhra Pradesh, India

A study on the tetracycline resistance *Staphylococcus aureus* from milk

K Harish, A Jagadeesh Babu, T Madhava Rao, B Somasekhara Goud and Y Suresh

Abstract

The present study was undertaken to isolate the *Staphylococcus aureus* from raw milk samples obtained from local vendors, to study their antibiotic sensitivity/resistant patterns, the molecular characterization of species specific *Staphylococcus aureus* and their methicillin resistance and tetracycline resistance by using the primers derived from *nuc* and *tetK* genes which gave specific amplification product of 270 bp and 360 bp for *S. aureus* and tetracycline resistance respectively. Out of 400 samples collected from different sources, 374 samples were positive for *Staphylococcus aureus*, Among the 374 isolates 191 isolates were confirmed as pathogenic *S. aureus* by a positive coagulase test. A pannel of 8 antibiotic discs were tested by using the standard disc diffusion method. Among the 191 isolates maximum resistance was observed for penicillin (74.9%) followed by gentamycin (27.2%), erythromycin (21%), cefoxitin (15.2%), and ciprofloxacin (11.5%), tetracycline (7.8%), cotrimoxazole (5.8%), chloramphenicol (0.5%). Out of 400 samples 374 samples were positive for *S. aureus* by cultural methods and 191 (51.07%, 191/374) were positive by PCR method. All the coagulase positive *S. aureus* isolates (191) were positive for *nuc* gene. In the present study for all the tetracycline resistant *S. aureus* isolates PCR assay was conducted by targeting *tetK* gene. Out of 15 tetracycline resistant *S. aureus* isolates 12 isolates were positive for *tetK* gene. This accounts for 6.3% of the total *S. aureus* isolates.

Keywords: *Staphylococcus aureus*, molecular characterization-antimicrobial resistance, *nuc* gene and *tetK* gene

Introduction

Staphylococcus aureus is a common microorganism found in raw milk and has been associated with food poisoning due to consumption of raw milk and contaminated dairy products and it also causes a wide variety of diseases from mild skin infection to more severe diseases such as pneumonia and septicaemia. Presence of *S. aureus* in milk can be due to poor hygiene practices of milk handlers because *S. aureus* is naturally present on the hands, nasal cavity and skin of human. Other than milk handlers and farm workers, the cow itself can be a source of *S. aureus*, especially if it is suffering from clinical or subclinical mastitis or having skin lesions. In Tirupati and nearby villages, milking is being done by hand and this might increase the risk of direct human contact with the raw milk. Contaminated milking equipments and the milker's hands also may be the source of infection. Foodborne infection risk is low in the countries where pasteurization is applied to most milk products, but there exists a risk with raw milk and products made with raw milk.

Antimicrobial agents are normally administered to cattle to treat microbial infections. Prolonged use of antimicrobial agents may lead to the emergence of antimicrobial resistant bacterial strains which is a serious concern not only in animal health but also more importantly to human health. Presence of antimicrobial resistant genes in *Staphylococcus* species is also of great concern since resistance genes can be transferred between staphylococcal species through lateral transfer and these pathogens harbouring resistant genes can be transferred to humans from animals. In *Staphylococcus* species some of the genes responsible for antibiotic resistance like *mecA* (methacillim), *blaZ* (penicillin) *tetM/tetK* (tetracycline), *ermA/ermC* have become public health concern.

The development of multiple antibiotic-resistant bacteria due to indiscriminate of antibiotics in animals and poultry production is well authenticated for pathogenic bacteria (Hawkey *et al.*, 2009; Arenas *et al.*, 2017) [19, 5]. Contaminated food of animal origins with antibiotic-resistant bacteria can be a great threat to public health, and the antibiotic resistance determinants can be transferred from antibiotic-resistant bacteria to other bacteria affecting human

(Threlfall *et al.*, 2000) [49]. Identical elements of antibiotic-resistant genes found in bacteria that affect both animals and humans have shown the role of raw foods in the dissemination of these resistance genes through the food chains (Teuber, 2001) [47] or through occupational contact with livestock (Leibler *et al.*, 2016) [26]. The multiple antibiotic-resistant bacteria were commonly isolated from food of animal origin such as raw milk and unpasteurized dairy products and meat products (Fawzy *et al.*, 2017) [15], the resistance genes can be transferred from antibiotic-resistant bacteria to the intestinal flora of humans through food products, and these bacteria can be a reservoir of resistant genes for pathogenic bacteria (Aarestrup *et al.*, 2008) [1]. The development of tetracycline resistance in *S. aureus* by acquisition of *tetK*, *tetL*, *tetM*, *tetO* genes. Plasmid encoded *tetK* gene was responsible for tetracycline resistance which mediates active efflux mechanism, and most often seen in *S. aureus* among

tetracycline resistant genes. The *tetK* gene confers resistance to tetracycline and chlortetracycline but neither to minocycline nor doxycycline.

Keeping in view of above reasons the present investigation was designed to study the antibiotic resistance profile of *Staphylococcus aureus* isolated from the milk samples collected in and around Tirupati.

Materials and Methods

For this study raw milk samples were collected by using sterile sampling tubes. A total of 400 milk samples were collected aseptically from different sources in and around Tirupati, Andhra Pradesh (table 1). The source of the milk samples comprises of individual farmers and local street vendors. The milk samples were processed within 2 to 24 hours of collection.

Table 1: Source and number of raw milk samples collected

Sl. No.	Place of sample collection	No. of samples
1.	Gurava Reddy Nagar	19
2.	Tummalagunta	16
3.	Sri Nagar	36
4.	Rajeev Nagar	32
5.	VK puram	12
6.	Lingeswara Nagar	54
7.	Padmavathi Nagar	52
8.	Peruru	120
9.	Cherlapalli	24
10.	Ambedkar Nagar	35
	Grand Total	400

Tryptic soy broth was used for enrichment of inoculums. Baird Parker agar supplemented with 5% sterile egg yolk tellurite suspension was used for isolation of *S. aureus*. 1ml of milk sample was inoculated in 9ml of sterile enrichment broth and incubated for 24 h at 37 °C. Subsequently, a loop full of inoculum from enrichment broth was streaked on Baird Parker agar (Hi Media Pvt. Ltd., India). After 48 h of incubation at 37 °C, colonies appeared as jet black surrounded by a white halo were identified as characteristic growth of *S. aureus*. The typical colonies were selected and the smears prepared were stained by Gram's staining. The presence of Gram-positive cocci and showing bunch of grapes like appearance were presumed as *Staphylococci* spp. For confirmation of *S. aureus*, the biochemical test conducted were catalase test, oxidase test, coagulase test, IMViC tests

and Triple sugar iron test.

The isolates were subjected to antibiotic resistance profile study against 8 antibiotics by using disk diffusion method according to Clinical and Laboratory Standards Institute (CLSI, 2016) guidelines. *S. aureus* isolates were inoculated into BHI broth and incubated at 35 °C for 24h. A bacterial suspension with the turbidity adjusted to a 0.5 McFarland standard was seeded on Muller Hinton agar plates and spread by the sterile cotton swabs. Then the plates were allowed to dry and antibiotic discs were placed aseptically with sterile fine forceps. The plates were incubated at 35 °C for 16-18h and Cefoxitin is kept for 24h incubation. The diameter of the zone of inhibition was compared with interpretive standards for *S. aureus* given by CLSI (Table- 2). After the incubation period was over inhibition zones were measured.

Table 2: Interpretation chart for antibiotic sensitivity/resistance patterns (Hi-Media)

S. No	Antimicrobial agent	Disc Conc.	Diameter of zone of inhibition in mm		
			Sensitive	Intermediate resistant	Resistant
1.	Pencillin-G (P)	10 units	≥29	-	≤28
2.	Cefoxitin (CX)	30mcg	≥22	-	≤21
3.	Tetracycline (TE)	30 mcg	≥19	15-18	≤14
4.	Cotrimoxazole (COT)	25mcg	≥16	11-15	≤10
5.	Erythromycin (E)	15 mcg	≥23	14-22	≤13
6.	Gentamicin (GEN)	10 mcg	≥15	13-14	≤12
7.	Chloramphenicol (C)	30 mcg	≥18	13-17	≤12
8.	Ciprofloxacin(CIP)	5mcg	≥21	16-20	≤15

The turbidity standard for 0.5 McFarland standard was prepared by adding 0.5 ml of (1.17% w/v) Barium chloride dehydrate (BaCl₂ 2H₂O) solution to 1% Sulphuric acid. The turbidity standard was placed in a tube identical to the one used for the broth sample and was stored in the dark at room

temperature. The turbidity was equivalent to 10⁸cfu/ml which is half the density of a Mac Farland 0.5 standard. The standard was agitated on a vortex mixer immediately before use. If the culture was found less turbid than the turbidity standard it was further incubated for 2-8 hours at 37 °C until

turbidity was equivalent to the standard. If the turbidity exceeds that of the standard the culture solution was diluted with tryptic soya broth to equitate with the standard.

After identification, biochemical characterization and detection of antibiotic resistance profile of the isolates the pure cultures were streaked on Nutrient agar slant and incubated for 24 h at 37 °C and then stored at 4 °C for further analysis.

Molecular characterization of the isolates for *nuc* gene and *tetK* gene was carried out by Polymerase Chain Reaction. The reference strains for *Staphylococcus aureus* (MTCC3103) and for Methicillin Resistant *Staphylococcus aureus* (ATCC 33591) were obtained from Department of Veterinary Public

Health & Epidemiology, College of Veterinary Science, Tirupati, Andhra Pradesh. For the extraction of DNA Suspensions of the bacterial colonies maintained on the nutrient agar slant were prepared in 1.5 ml microcentrifuge tubes in 250 µl of sterile double distilled water by gentle mixing. The samples were boiled for 10 min, cooled on ice for 10 min and centrifuged at 10,000rpm for 10 min. 4µl of the supernatant was used as the template for each polymerase chain reaction (PCR) (Islam *et al.*, 2016) [21]. The primers used in the study were custom synthesized by Eurofins genomics India Private Limited, Bangalore. The details of the primers are given in Table. 3.

Table 3: Details of oligonucleotide primers used in this study

Target gene	Primer sequence (5 ¹ -3 ¹)	Amplicon size (bp)	Reference
<i>nuc</i>	F- GCG ATT GAT GGT GAT ACG GTT	270	Bharathy <i>et al.</i> (2015) [7]
	R- AGC CAA GCC TTG ACG AAC TAA AGC		
<i>tetK</i>	F- GTA GCG ACA ATA GGT AAT AGT	360	Arafa <i>et al.</i> (2016) [4]
	R- GTA GTG ACA ATA AAC CTC CTA		

Initial experiments to optimize PCR reaction conditions for *S. aureus* template involved the empirical variation of annealing temperature (53 °C – 56 °C), concentration of primer (5 – 15 pmol), template volume (2µl - 5µl) and the cycling conditions. Optimal results were obtained using 5 µl of bacterial lysate or 20ng of diluted DNA as template in a master mix (2X). Amplification for *nuc* gene was carried by 5 min initial denaturation at 94 °C followed by 35 cycles of denaturation at 94 °C for 1min, annealing at 55 °C for 40sec and extension at 72 °C for 1 min and a final extension for 10 min at 72 °C. Amplification of *tetK* gene was carried by 5 min initial denaturation at 94 °C followed by 35 cycles of denaturation at 94 °C for 30sec, annealing at 54 °C for 30sec and extension at 72 °C for 30sec and a final extension for 10 min at 72 °C. The given PCR conditions was found to be optimum for obtaining the desired PCR amplification of 270 bp from *nuc* gene and 360 bp from *tetK* gene of *S. aureus*.

A simplex PCR assay was developed for the detection of species specific gene of *S. aureus* (*nuc* gene) and gene responsible for tetracycline resistance of *S. aureus* (*tetK* gene) was used in our study according to Bharathy *et al.* (2015) [7] and Arafa *et al.* (2016) [4] with slight modifications. The simplex PCR protocol for each gene was followed for the detection of various genes using standard protocols. Following initial trials with varying concentrations of components, the reaction mixture for each gene was optimized as below indicated in the following tables-4&5.

Table 6: Cyclic conditions used for simplex PCR assay targeting *nuc* gene and *tetK* gene.

Gene	Denaturation		Annaeling		Extension	
	Tempeature (°C)	Time (Seconds)	Tempeature (°C)	Time (seconds)	Tempeature (°C)	Time (seconds)
<i>nuc</i>	94	60	55	40	72	60
<i>tetK</i>	94	30	54	30	72	30

The resultant PCR products were subjected to 1.5% agarose gel electrophoresis as described by Sambrook and Russel (2001) [42]. Agarose gel (1.5%) was prepared by boiling 0.6 gm of agarose in 40 ml of 1X TBE buffer. After cooling, ethidium bromide was added to the agarose solution to a final concentration of 0.5µg / ml. The molten agarose was poured into a gel casting tray fitted with acrylic comb was kept undisturbed till the gel has solidified. After solidification of the gel, a few ml of 1X TBE buffer was added and the comb

Table 4: Optimized reaction mixture for detection of *nuc* gene

Name	µl per reaction
PCR Master mix(2X)	12.5
<i>nuc</i> gene(Forward primer) (10pmol)	1.0
<i>nuc</i> gene(Reverse primer) (10pmol)	1.0
Template DNA	4.0
nuclease free water	6.5
Total	25.0

Table 5: Optimized reaction mixture for detection of *tetK* gene

Name	µl per reaction
PCR Master mix(2X)	12.5
<i>tetK</i> gene (Forward primer) (10pmol)	1.0
<i>tetK</i> gene (Reverse primer) (10pmol)	1.0
Template DNA	4.0
nuclease free water	6.5
Total	25.0

DNA amplification was performed in a Thermal cycler with a pre-heated lid. The cycling conditions for detection of all genes were depicted in the table-6. The resultant PCR products were stored at 4 °C in TE buffer for further analysis by 1.5% agarose gel electrophoresis stained with ethidium bromide. PCR was run for 35 cycles with initial denaturation at 94 °C for 5 minutes and final extension at 72 °C for 10 min for all the oligonucleotide primer sets. (Bharathy *et al.*, 2015 and Arafa *et al.*, 2016) [4, 7]

was removed carefully and then the tray containing the gel was placed in a submarine horizontal electrophoresis unit filled with 1X TBE buffer up to a level of 1mm above the gel surface.

About 5µl of each PCR product was mixed with 1µl of gel loading dye (6X) and loaded into each well. Electrophoresis was performed at 5V/cm and the mobility was monitored by the migration of the dye. After sufficient migration, the gel was observed under UV transillumination using Alpha

innotech gel documentation system to visualize the bands. The PCR product size was determined by comparing with a standard molecular weight marker.

Results & Discussion

In the present investigation, a total of 400 raw milk samples were collected in and around Tirupati city, Andhra Pradesh. All the milk samples were subjected to isolation and identification by conventional methods and from a total of 400 milk samples 374 (93.5%) samples were positive for staphylococcal species by culture methods. For confirmation of *S. aureus*, all the 374 isolates were subjected to different biochemical tests like catalase test, oxidase test, indole test, methyl red test, Voges Proskauer test, citrate utilization test and triple sugar iron agar test. All the 374 isolates were found catalase positive, oxidase negative, indole test negative, methyl red test positive, Voges Proskauer test positive, citrate utilization test negative and triple sugar iron test positive by yellow discolouration of slant and butt without production of H₂S gas. The present findings were in agreement with Reddy *et al.* (2015) [40], Habib *et al.* (2015) [17] and Ramya *et al.* (2017) [39] who have subjected the isolates of *staphylococci* to same biochemical tests and found similar type of reactions.

Multi drug resistance is now the norm among the Gram Positive bacteria like pneumococci, enterococci and staphylococci. *S. aureus* is perhaps the pathogen of concern because of its intrinsic virulence, its ability to cause a diverse array of life threatening infections in humans and in various animal species and its capacity to adapt to different environmental conditions. The evolution of increasingly antimicrobial resistant bacteria stems from a multitude of factors including the strong selective pressure caused by the wide spread and sometimes inappropriate use of antimicrobial agents. The increase in regional and international travel and the relative ease with which antimicrobial resistant bacteria cross geographic barriers (Lowy, 2003)

During the present study, all *S. aureus* isolates were found variably resistant to the antibiotics tested. Among the 191 *S. aureus* isolates maximum resistance was observed for penicillin-G (74.9%) followed by gentamycin (27.2%), erythromycin (21%), cefoxitin (15.2%), ciprofloxacin (11.5%), tetracycline (7.8%), cotrimoxazole (5.8%) and chloramphenicol (0.5%).

In the present study, *S. aureus* isolates have showed highest resistance to penicillin (74.9%) which was in agreement with the reports of Hanson *et al.* (2011) [18] and Fawzy *et al.* (2017) [15] who have found 77.7% and 73.6% resistance respectively. The findings of the present study with regarding the resistance of *S. aureus* to penicillin differ from earlier report of Khakpoor *et al.* (2011) [24], Thaker *et al.* (2013) [48], and Jahan *et al.* (2015) [23] who have recorded 100% resistance to penicillin among *S. aureus* isolates. Slightly higher percentage of resistance to penicillin than the findings of this investigation was observed from the findings of Abera *et al.* (2010) [2], Tigabu *et al.* (2015) [50], Awad *et al.* (2017) [6], Feng *et al.* (2016) [16], Elemo *et al.* (2017) [13], Can *et al.* (2017) and Yadav, (2018) [54], who have reported 94.4%, 83.5%, 83.3%, 84.09%, 87.3%, 81.81% and 82.23% of resistance respectively.

Resistance to cefoxitin was found 15.2% in *S. aureus* isolates of the present study which was in agreement with the report of Shamila-Syuhada *et al.* (2016) [44] who found 15% resistance in Penang, Malaysia. Higher resistance to cefoxitin by the isolates of *S. aureus* compare to the present finding

was observed in Tigabu *et al.* (2015) [50], Özdemir and Keyvan *et al.* (2016) [36], Ammar *et al.* (2016) [3], and Elemo *et al.* (2017) [13] who have reported 53.2%, 22.8%, 55%, and 58.1% of resistance respectively. In contrast to the present findings lower resistance to cefoxitin by *S. aureus* was observed by Mashouf *et al.* (2015) [30] and Can *et al.* (2017) who have observed 5.1% and 9.09% of resistance respectively.

In the present investigation resistance to tetracycline was found to be a 7.8% among the isolates of *S. aureus* which was in agreement with the report of Shamila-Syuhada *et al.* (2016) [44] who found 5% resistance to tetracycline among the isolates of *S. aureus* in Penang, Malaysia. Higher resistance compared to the present finding was observed for tetracycline resistance among the isolates of *S. aureus* in the findings of Mirzaei *et al.* (2012), Mubarack *et al.* (2012) [35], Jackson *et al.* (2013) [22], and Feng *et al.* (2016) [16] who have reported 23%, 11.84%, 25%, and 15.91% of resistance respectively. Only 1% of resistance for tetracycline was observed among the isolates of *S. aureus* in the findings of Wang *et al.* (2018) [53] which was found to be lower than the findings of the present investigation.

Resistance to gentamicin in the present study by the isolates of *S. aureus* was 27.2% which was in agreement with the reports of Momtaz *et al.* (2013) [34] and Mashouf *et al.* (2015) [30] who found 29.26% and 27.6% of resistance in the isolates of *S. aureus* respectively. Higher resistance compared to the present findings was observed by Jackson *et al.* (2013) [22] who reported 33.3% resistance in Georgia. Lower resistance to gentamycin compared to the present findings was observed by Thaker *et al.* (2013) [48], Tigabu *et al.* (2015) [50], Feng *et al.* (2016) [16] and Wang *et al.* (2018) [53] who have reported 10%, 2.8%, 9.09% and 1% respectively.

The isolates of *S. aureus* in present study have shown 21% of resistance to erythromycin which was in agreement with the report of Feng *et al.* (2016) [16] who found 20.45% resistance in Northwest China. Higher resistance to erythromycin by the isolates of *S. aureus* compared to the present finding was observed in Mashouf *et al.* (2015) [30] and Fawzy *et al.* (2017) [15] who reported 30.6% and 26.3% resistance respectively.

Lower resistance to erythromycin compare to the present finding was observed by Hanson *et al.* (2011) [18], Tigabu *et al.* (2015) [50], Can *et al.* (2017), Hoque *et al.* (2018) [23] and Wang *et al.* (2018) [53] who have observed 14.8%, 4.6%, 16%, 8.2% and 5.2% of resistance respectively.

Resistance to cotrimoxazole was found to be a 5.8% by the isolates of *S. aureus* in the present study which was in agreement with the report of Tigabu *et al.* (2015) [50] who found 4.6% resistance among the isolates of *S. aureus* in central highlands of Ethiopia. Higher resistance compared to the present finding towards the cotrimoxazole by the isolates of *S. aureus* was observed in the findings of Mashouf *et al.* (2015) [30] and Pourtaghi *et al.* (2015) who have reported 14.3% and 11.18% resistance respectively in Iran. Lower resistance to cotrimoxazole than the present findings was observed by Mirzaei *et al.* (2012), Özdemir and Keyvan *et al.* (2016) [36] and Ammar *et al.* (2016) [3] who have observed 3%, 3.5%, and 2.5% resistance respectively.

The isolates of *S. aureus* have shown resistance to ciprofloxacin and it was found as 11.5% which was in agreement with the report of Ammar *et al.* (2016) [3] and Awad *et al.* (2017) [6] who have observed 10% and 14.3% resistance in the isolates of *S. aureus* in Egypt. Higher resistance compare to the present finding was observed by Kumar *et al.* (2011), Kreausukon *et al.* (2012) [25], Pati *et al.*

(2016)^[37], Wang *et al.* (2018)^[53] and Yadav, (2018)^[54] who have reported 26.2%, 24.5%, 37%, and 18.8% and 42.1% of resistance respectively among the isolates of *S. aureus*. Lower resistance to ciprofloxacin compared to the present findings was observed in the findings of Tigabu *et al.* (2015)^[50] who reported 3.7% of resistance in his study.

Resistance to chloramphenicol by the isolates of *S. aureus* in the present investigation was found only 0.5% which was in agreement with the report of Wang *et al.* (2018)^[53] who found 1% resistance in his isolates of *S. aureus* in Beijing, China. Higher resistance compared to the present finding was observed in the findings of Ammar *et al.* (2016)^[3], Pati *et al.* (2016)^[37] and Özdemir and Keyvan *et al.* (2016)^[36] who have reported 12.5%, 9% and 5.2% resistance respectively. No resistant isolates for chloramphenicol were observed in the findings of Abera *et al.* (2010)^[2], Dittaman *et al.* (2017), Elemo *et al.* (2017)^[13]. Due to reduced usage of chloramphenicol in the field condition, negligible amount of resistance was observed in the present study. Although it is extremely difficult to explain these conflicting data with regards to both time and place of study, the variation is probably due to differential clonal expression and drug pressure in community.

Several workers have used PCR with varied success for detection of MRSA from clinical samples using specific gene primers for targeting. Of the specific gene sequences *nuc* has been most frequently targeted for PCR based detection of *staphylococcus*. The *nuc* gene has been designated as species specific gene for *Staphylococcus aureus*, because *S. aureus* strains produce an extra cellular thermostable nuclease (thermonuclease, TNase) with a frequency similar to that as which they produce coagulase enzyme (Madison *et al.*, 1983)^[28]. The TNase protein has been well characterized and its gene the *nuc* gene has been cloned and sequenced (Tucker *et al.*, 1978)^[52]. Thus the *nuc* gene which encodes the thermostable nuclease that is highly specific for *S. aureus* (Saha *et al.*, 2008)^[41].

The isolates positive for *S. aureus* by culturing and biochemical tests were further confirmed by PCR targeting for species specific gene of *S. aureus* (*nuc* gene). *nuc* gene codes for thermostable nuclease (TNase) which is specific for *S. aureus* (Bharathy *et al.*, 2015)^[7].

In the present study, out of 400 samples screened, 47.75% (191/400) *S. aureus* isolates were positive for *nuc* gene. This was nearly in agreement with findings of Medvedová *et al.* (2014) and Awad *et al.* (2017)^[6] who found 44% and 42% of the isolates were *S. aureus* respectively. Higher incidence compared to the present findings was reported by Cremonesi *et al.* (2007)^[11], Falaki and Mahdavi *et al.* (2017)^[14], Pati *et al.* (2016)^[37] and Bharathy *et al.* (2015)^[7] who have found 100%, 85%, 83.8% and 65.57% of *S. aureus* isolates from their study and were characterized by *nuc* gene. Lower incidence of *nuc* gene compared to the present findings was reported by Tanzin *et al.* (2016)^[46], Srednik *et al.* (2018)^[45] and Mashouf *et al.* (2015)^[30] who have reported 35.29%, 28.1% and 23.52% of *S. aureus* isolates were *nuc* gene respectively.

Plasmid encoded *tetK* gene was responsible for tetracycline resistance which mediates active efflux mechanism, and most oftenly seen in *S. aureus* among tetracycline resistant genes (Trzcinski *et al.*, 2000; Chopra *et al.*, 2001)^[51, 9]. The *tetK* gene confers resistance to tetracycline and chlortetracycline but neither to minocycline nor doxycycline.

In the present study, for all the tetracycline resistant *S. aureus* isolates PCR assay was conducted for targeting *tetK* gene. Out of 15 tetracycline resistant *S. aureus* isolates 12 isolates were positive for *tetK* gene. This accounts for 6.3% of the total *S. aureus* isolates. This was moderately in agreement with the finding of Sharma *et al.* (2017)^[43] who found *tetK* gene in 15% of his isolates of *S. aureus*. Higher percentage of *tetK* gene compared to the present findings was observed by Arafa *et al.* (2016)^[4], Fawzy *et al.* (2017)^[15], Martini *et al.* (2017), Memon *et al.* (2013)^[32] and Momtaz *et al.*, (2013)^[34] who have found that 56%, 52.43%, 100%, 100%, and 84% of their isolates of *S. aureus* have shown *tetK* gene respectively. While lower incidence compared to the present finding was observed in Shamila-Syuhada *et al.* (2016)^[44] who found only 1.67% of isolates have shown *tetK* gene.

Antibiogram Representation by Graphs




Fig 1: *S. aureus* showing sensitivity patterns to different antibiotics




Fig 2: *S. aureus* showing intermediate resistance patterns to different antibiotics




Fig 3: *S. aureus* showing resistance patterns to different antibiotics




Fig 4: *S. aureus* isolates showing *nuc* gene

Lane M	Molecular marker(100bp)
Lane 2, 3, 4, 5 and 6	<i>S. aureus</i> isolates carrying <i>nuc</i> gene(270bp)




Fig 5: *S. aureus* isolates showing *tetK* gene

Lane M	Molecular marker(100bp)
Lane 2, 3, 4, 5 and 6	<i>S. aureus</i> isolates carrying <i>tetK</i> gene(360bp)

Conclusion

The present investigation revealed that there is an increased trend in the presence of antibiotic resistant *Staphylococcus aureus* in the milk samples collected from the vendors. As the antimicrobial resistant microflora is having greatest public health threat, necessary precautions have to be taken with regarding the administration of antibiotics through feed and indiscriminate use of antibiotics in the treatment should also be curtailed. Bringing awareness among the public about the harmful effects of multi drug resistant microflora is another important objective for the scientists to protect the humans from these super bugs.

Acknowledgement

The present study is the M.V.Sc. thesis work of Dr. K. Harish and the authors. Thank Sri Venkateswara Veterinary University, Tirupati, Andhra Pradesh, for providing necessary facilities and funds to carry out the work.

References

1. Aarestrup FM, Wegener HC, Collignon P. Resistance in bacteria of the food chain: epidemiology and control strategies. *Expert review of anti-infective therapy*. 2008; 6(5):733-750.
2. Abera M, Demie B, Aragaw K, Regassa F, Regassa A. Isolation and identification of *Staphylococcus aureus* from bovine mastitic milk and their drug resistance patterns in Adama town, Ethiopia. *Journal of Veterinary*

3. Ammar AM, Attia AM, Abd El-Hamid MI, El-Shorbagy IM, Abd El-Kader SA. Genetic basis of resistance waves among methicillin resistant *Staphylococcus aureus* isolates recovered from milk and meat products in Egypt. *Cellular and Molecular Biology (Noisy-le-Grand)*. 2016; 62(10):7-15.
4. Arafa AA, Ibrahim ES, Fouad EA, Gaber ES. Antibiotic resistance of *Staphylococci* concerning strains included in food industry in Egypt. *International Journal of Pharmaceutical and Clinical Research*. 2016; 8(12):1583-1589.
5. Arenas NE, Abril DA, Valencia P, Khandige S, Soto CY, Moreno-Melo V. Screening foodborne and zoonotic pathogens associated with livestock practices in the Sumapaz region, Cundinamarca, Colombia. *Tropical animal health and production*. 2017; 49(4):739-745.
6. Awad A, Ramadan H, Nasr S, Ateya A, Atwa S. Genetic characterization antimicrobial resistance patterns and virulence determinants of *Staphylococcus aureus* isolated from bovine mastitis. *Pakistan Journal of Biological Sciences*. 2017; 20(6):298-305.
7. Bharathy S, Gunaseelan L, Porteen K, Bojiraj M. Prevalence of *Staphylococcus aureus* in raw milk: can it be a potential public health threat? *International Journal of Advanced Research*. 2015; 3(2):801-806.
8. Can HY, Elmalı M, Karagöz A. Molecular typing and antimicrobial susceptibility of *Staphylococcus aureus* strains isolated from raw milk, cheese, minced meat, and chicken meat samples. *Korean journal for food science of animal resources*. 2017; 37(2):175-180.
9. Chopra I, Roberts M. Tetracycline antibiotics: mode of action, applications, molecular biology, and epidemiology of bacterial resistance. *Microbiology and molecular biology reviews*. 2001; 65(2):232-260.
10. CLSI. "Performance standards for antimicrobial susceptibility testing". 26th Informational supplement M100-S, Clinical and Laboratory Standard Institute, Wayne, Pennsylvania, USA. 2016.
11. Cremonesi P, Perez G, Pisoni G, Moroni P, Morandi S, Luzzana M *et al*. Detection of enterotoxigenic *Staphylococcus aureus* isolates in raw milk cheese. *The Society for Applied Microbiology, Letters in Applied Microbiology*. 2007; 45(6):586-591.
12. Dittmann KK, Chaul LT, Lee SH, Corassin CH, Fernandes de Oliveira CA., Pereira De Martinis EC *et al*. *Staphylococcus aureus* in some Brazilian dairy industries: changes of contamination and diversity. *Frontiers in Microbiology*. 2017; 8:2049.
13. Elemo KK, Sisay T, Shiferaw A, Fato MA. Prevalence, risk factors and multidrug resistance profile of *Staphylococcus aureus* isolated from bovine mastitis in selected dairy farms in and around Asella town, Arsi Zone, South Eastern Ethiopia. *African Journal of Microbiology Research*. 2017; 11(45):1632-1642.
14. Falaki B, Mahdavi S. Study of distribution of biofilm producing genes in *Staphylococcus aureus* isolated from local cheese samples in Maragheh city, Iran. *Gene Cell Tissue (In Press)*. 2017; 4(4):1-5.
15. Fawzy R, Samy AA, Salam HS, Khairy EA, Koraney AA. Polymerase chain reaction detection of genes responsible for multiple antibiotic resistance *Staphylococcus aureus* isolated from food of animal origin in Egypt. *Veterinary world*. 2017; 10(10):1205-

- 1211.
16. Feng Y, Qi W, Wang XR, Ling W, Li XP, Luo JY *et al.* Genetic characterization of antimicrobial resistance in *Staphylococcus aureus* isolated from bovine mastitis cases in Northwest China. *Journal of integrative agriculture*. 2016; 15(12):2842-2847.
 17. Habib F, Rind R, Durani N, Bhatto AL, Buriro RS, Tunio A *et al.* Morphological and cultural characterization of *Staphylococcus aureus* isolated from different animal species. *Journal of Applied Environmental and Biological Sciences*. 2015; 5(2):15-26.
 18. Hanson BM, Dressler AE, Harper AL, Scheibel RP, Wardyn SE, Roberts LK *et al.* Prevalence of *Staphylococcus aureus* and methicillin resistant *Staphylococcus aureus* (MRSA) on retail meat in Iowa. *Journal of infection and public health*. 2011; 4(4):169-174.
 19. Hawkey PM, Jones AM. The changing epidemiology of resistance. *Journal of Antimicrobial Chemotherapy*. 2009; 64(1):3-10.
 20. Hoque MN, Das ZC, Rahman ANMA, Haider MG, Islam MA. Molecular characterization of *Staphylococcus aureus* strains in bovine mastitis milk in Bangladesh. *International Journal of Veterinary Science and Medicine*. 2018; 6(1):53-60.
 21. Islam MA, Kabir SML, Rahman MT. Molecular detection and characterization of *Staphylococcus aureus* isolated from raw milk sold in different markets of Bangladesh. *Bangladesh Journal of Veterinary Medicine*. 2016; 14(2):277-282.
 22. Jackson CR, Davis JA, Barrett JB. Prevalence and characterization of methicillin resistant *Staphylococcus aureus* isolates from retail meat and humans in Georgia. *Journal of Clinical Microbiology*. 2013; 51(4):1199-1207.
 23. Jahan M, Rahman M, Parvej MS, Chowdhury SMZH, Haque ME, Talukder MAK *et al.* Isolation and characterization of *Staphylococcus aureus* from raw cow milk in Bangladesh. *Journal of Advanced Veterinary and Animal Research*. 2015; 2(1):49-55.
 24. Khakpoor M, Safarmashaei S, Jafary R. Study of milk extracted from cows related to *Staphylococcus aureus* by culturing and PCR. *Global Veterinaria*. 2011; 7(6):572-575.
 25. Kreausakon K, Fetsch A, Kraushaar B, Alt K, Müller K, Krömker V *et al.* Prevalence, antimicrobial resistance, and molecular characterization of methicillin resistant *Staphylococcus aureus* from bulk tank milk of dairy herds. *Journal of Dairy Science*. 2012; 95(8):4382-4388.
 26. Leibler JH, Jordan JA, Brownstein K, Lander L, Price LB, Perry MJ. *Staphylococcus aureus* nasal carriage among beef packing workers in a Midwestern United States Slaughter house. *PLoS one*. 2016; 11(2):1-11.
 27. Lowy FD. *Staphylococcus aureus* infections. *The New England Journal of Medicine*. 1998; 339(8): 520-532.
 28. Madison BM, Baselski VS. Rapid identification of *Staphylococcus aureus* in blood cultures by thermo nuclease testing. *Journal of clinical microbiology*. 1983; 18(3):722-724.
 29. Martini CL, Lange CC, Brito MA, Ribeiro JB, Mendonça LC, Vaz EK. Characterisation of penicillin and tetracycline resistance in *Staphylococcus aureus* isolated from bovine milk samples in Minas Gerais, Brazil. *Journal of Dairy Research*. 2017; 84(2):202-205.
 30. Mashouf RY, Hosseini SM, Mousavi SM, Arabestani MR. Prevalence of enterotoxin genes and antibacterial susceptibility pattern of *Staphylococcus aureus* strains isolated from animal originated foods in west of Iran. *Oman Medical Journal*. 2015; 30(4):283-290.
 31. Medvedová A, Studenicova A, Valik L, Horvathova Z. Prevalence and growth dynamics of enterotoxigenic *Staphylococcus aureus* isolates in Slovakian dairy products. *Czech Journal of Food Science*. 2014; 32(4):337-341.
 32. Memon J, Yang Y, Kashif J, Yaqoob M, Buriro R, Soomro J *et al.* Genotypes, virulence factors and antimicrobial resistance genes of *Staphylococcus aureus* isolated in bovine subclinical mastitis from Eastern China. *Pakistan Veterinary Journal*. 2013; 33(4):486-491.
 33. Mirzaei H, Farhoudi H, Tavassoli H, Farajli M, Monadi A. Presence and antimicrobial susceptibility of methicillin resistant *Staphylococcus aureus* in raw and pasteurized milk and ice cream in Tabriz by culture and PCR techniques. *African Journal of Microbiology Research*. 2002; 6(32):6224-6229.
 34. Momtaz H, Dehkordi FS, Rahimi E, Asgarifar A, Momeni M. Virulence genes and antimicrobial resistance profiles of *Staphylococcus aureus* isolated from chicken meat in Isfahan province, Iran. *Journal of Applied Poultry Research*. 2013; 22(4):913-921.
 35. Mubarack HM, Doss A, Vijayasanthi M, Venkataswamy R. Antimicrobial drug susceptibility of *Staphylococcus aureus* from subclinical bovine mastitis in Coimbatore, Tamilnadu, South India. *Veterinary World*. 2012; 5(6):352-355.
 36. Özdemir H, Keyvan E. Isolation and characterisation of *Staphylococcus aureus* strains isolated from beef, sheep and chicken meat. *Ankara Üniversitesi Veteriner Fakültesi Dergisi*. 2016; 63:333-338.
 37. Pati BK, Mukherjee R. Characterization of *Staphylococcus aureus* isolates of bovine mastitis origin and antibiotic sensitivity pattern from Northern plains of India. *Journal of Veterinary Research and Animal Husbandry*. 2016; 1(1):1-5.
 38. Pourtaghi H, Azizi AG, Sodagari H. Antimicrobial resistance patterns of *Staphylococcus aureus* isolated from bovine subclinical mastitis in Alborz province, Iran. *Bulgarian Journal of Veterinary Medicine*. 2016; 19(2):169-174.
 39. Ramya P. Prevalence of *Staphylococcus aureus* in raw milk samples. *International Journal of Chemical Studies*. 2017; 5(5):1351-1353.
 40. Reddy MS, Babu AJ, Ramya P, Swetha CS. Molecular characterization of methicillin resistant *Staphylococcus aureus* from goats, pigs and their handlers. *International Journal of Microbiology Research*. 2015; 7(3):648-655.
 41. Saha B, Singh AK, Ghosh A, Bal M. Identification and characterization of a vancomycin-resistant *Staphylococcus aureus* isolated from Kolkata (South Asia). *Journal of medical microbiology*. 2008; 57(1):72-79.
 42. Sambrook J, Russell DW. *Molecular cloning: a laboratory manual*, 3rd edn. (Cold Spring Harbor Laboratory Press: New York), 2001.
 43. Sharma S, Bhowmik D, Bhattacharjee A. Tigecycline Resistance among Clinical Isolates of *Staphylococcus aureus* from North-east India. *Journal of Microbiology and Infectious Diseases*. 2017; 7(4):173-177.

44. Shamila-Syuhada AK, Rusul G, Wan-Nadiah WA, Chuah LO. Prevalence and antibiotics resistance of *Staphylococcus aureus* isolates isolated from raw milk obtained from small-scale dairy farms in Penang, Malaysia. *Pakistan Veterinary Journal*. 2016; 36(1):98-102.
45. Srednik ME, Usongo V, Lépine S, Janvier X, Archambault M, Gentilini ER. Characterisation of *Staphylococcus aureus* strains isolated from mastitis bovine milk in Argentina. *Journal of Dairy Research*. 2018; 85(1):57-63.
46. Tanzin T, Nazir KHMNH, Zahan MN, Parvej MS, Zesmin K, Rahman MT. Antibiotic resistance profile of bacteria isolated from raw milk samples of cattle and buffaloes. *Journal of Advanced Veterinary and Animal Research*. 2016; 3(1): 62-67.
47. Teuber M. Veterinary use and antibiotic resistance. *Current opinion in microbiology*. 2001; 4(5):493-499.
48. Thaker HC, Brahmabhatt MN, Nayak JB. Isolation and identification of *Staphylococcus aureus* from milk and milk products and their drug resistance patterns in Anand, Gujarat. *Veterinary World*. 2013; 6(1):10-13.
49. Threlfall EJ, Ward LR, Frost JA, Willshaw GA. The emergence and spread of antibiotic resistance in food-borne bacteria. *International journal of food microbiology*. 2000; 62(1-2):1-5.
50. Tigabu E, Kassa T, Asrat D, Alemayehu H, Sinmegn T, Adkins PRF *et al.* Phenotypic and genotypic characterization of *Staphylococcus aureus* isolates recovered from bovine milk in central highlands of Ethiopia. *African Journal of Microbiology Research*. 2015; 9(44):2209-2217.
51. Trzcinski K, Cooper BS, Hryniewicz W, Dowson CG. Expression of resistance to tetracyclines in strains of methicillin resistant *Staphylococcus aureus*. *Journal of Antimicrobial Chemotherapy*. 2000; 45(6):763-770.
52. Tucker PW, Hazen EE, Cotten FA. Staphylococcal nuclease reviewed: A prototypic study in contemporary enzymology, isolation, physical and enzymatic properties. *Mol. Cell Biochem*. 1978; 22:67-77.
53. Wang W, Lin X, Jiang T, Peng Z, Xu J, Yi L *et al.* Prevalence and characterization of *Staphylococcus aureus* cultured from raw milk taken from dairy cows with mastitis in Beijing, China. *Frontiers in microbiology*. 2018; 9(1123):1-16.
54. Yadav MM. Prevalence of *Staphylococcus aureus* in lactating cows with subclinical mastitis and their antibiogram in organized dairy farm, Maharashtra, India. *International Journal of Current Microbiology and Applied Sciences*. 2018; 7(3): 3674-3680.