



ISSN (E): 2277- 7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2021; 10(12): 1165-1168
© 2021 TPI
www.thepharmajournal.com
Received: 04-09-2021
Accepted: 10-10-2021

Lolamwad NS
M.Sc. Agriculture Scholar,
Department of Agronomy,
RCSM College of Agriculture,
Kolhapur, Maharashtra, India

Bhosale AS
Associate Professor, Department
of Agronomy, RCSM College of
Agriculture, Kolhapur,
Maharashtra, India

Solanke BN
M.Sc. Agriculture Scholar,
Department of Agronomy,
RCSM College of Agriculture,
Kolhapur, Maharashtra, India

Shinde RH
Associate Professor, Department
of Agronomy, RCSM College of
Agriculture, Kolhapur,
Maharashtra, India

Corresponding Author:
Lolamwad NS
M.Sc. Agriculture Scholar,
Department of Agronomy,
RCSM College of Agriculture,
Kolhapur, Maharashtra, India

Effect of fertilizer levels and plant densities on growth attribute and yield of summer sunflower (*Helianthus annuus* L.)

Lolamwad NS, Bhosale AS, Solanke BN and Shinde RH

Abstract

An present field experiment was conducted at the post graduate Research farm, Agronomy section, RCSM College of Agriculture, Kolhapur (MH), India. During March to April 2020 to investigate the effect of fertilizer levels and plant densities on growth attribute and yield of summer sunflower (*Helianthus annuus* L.). The experiment comprised three fertilizer levels viz, 45:45:45, 60:60:60 and 75:75:75 NPK kg ha⁻¹ and three plant densities viz, 30 cm x 30 cm, 45 cm x 30 cm and 60 cm x 30 cm. The experiment was laid out in Factorial randomized block design with four replication. Result revealed that fertilizer levels and plant densities had significant effect on growth attribute and yield of summer sunflower. The growth character and yield viz, plant height ⁻¹(cm), number of functional leaves⁻¹, leaf area (dm²), dry matter ⁻¹(g), seed yield, stalk yield and biological yield (q ha⁻¹) was recorded higher by application of fertilizer levels 75:75:75 kg NPK ha⁻¹. All the character were recorded higher under plant density (60 cm x 30 cm) except plant height, stalk yield and biological yield. The highest plant height, stalk yield and biological yield was recorded in (45 cm x 30 cm).

Keywords: Fertilizer levels, planting densities, sunflower, growth attributes, yield

Introduction

Sunflower (*Helianthus annuus* L.) is promising oil seed crop because of its short duration, photo insensitivity and wild adaptability to different agro-climatic regions and soil types. Sunflower can play an important role in meeting out the shortage of edible oil in country. Our country is facing acute shortage of edible oil mainly because of huge demand due to population pressure, raised standard of living and high demand from oil consuming industries. This demand is partly met by import of edible oils. Under such situation it needs to build up self-sufficiency in oil production and to meet the increasing demand of consumers. Sunflower oil is considered as a premium when compared to other vegetable oils. Sunflower oil is preferred among the consumers across world due to its health appeal and in India too, sunflower oil is the largest selling oil in the branded oil segment. Sunflower is also a crop of choice for farmers due to its wider adaptability, high yield potential, shorter duration and profitability. In India, the area under oilseed crop is 285.25 lakh ha with production of 328.77 lakh tones and productivity is 1094 kg ha⁻¹. The area under sunflower crop in world is 26.36 million hectares and production is 54.57 million tones with the productivity of 2090 kg ha⁻¹. Whereas, in India the area under sunflower crop is 0.23 million ha and production was, 0.17 million tones with the productivity of 760 kg ha⁻¹. In Maharashtra, sunflower is cultivated on an area of 25.1 thousand ha and production is 11.9 thousand tones with the productivity of 475.2 kg ha⁻¹ (Anonymous 2019) [3].

Plant densities affect growth characters and yield of sunflower through planting optimum LAI to reduce the competition between plants by more light penetration on plant canopies, consequently enhances photosynthesis process, then increase photosynthesis rate which consequently increase yield per unit area. Moreover, it is usually used in matching crop requirements to the environmental resources available. Furthermore, it is critical practices for determining the most favorable above ground surroundings that allows the plant to obtain the necessary environmental growth factors i.e. light, space, water, CO₂, and nutrients, which are significant tools to optimize the plant growth and accumulate and production of dry matter as well as consequently maximize the final seeds yield and its quality (Basha, 2000; Beg *et al.*, 2007 and Biabani, 2010) [4, 6]. In addition, number of plants per unit area play vital role in adjusting the crop environment and assist to improve disease avoidance, therefore modulating

number of plant per unit area is the main instrument to improve plant growth and the time required for plant canopy closure and to produce the highest quantity of biomass and seed yield (Khan *et al.*, 2003 and Liu *et al.*, 2008) [9, 12]. Long *et al.* (2001) [13] reported that due to the increasing of interplant competition for space, light, CO₂, water, and nutrients and other life factors, the growth characters and yield and its attributes of sunflower plants are probable to decline with raising the number of plants per unit area. While, under the optimum number of plants per unit area, growth, yield, and its attributes were expected to increase. Ali *et al.* (2012) [11] and Ibrahim (2012) [10] declared that plant growth characters, yield and attributes of sunflower plants significantly influenced by plant densities.

Nitrogen is essential constituents protein and chlorophyll, which is present in many other compounds of great physiological importance in plant metabolism such as nucleotides, phospholipids, alkaloids, enzyme, hormones, vitamins etc. it is thus very basic constituents of life. It important dark green colour to plant. Phosphorous plays important role in growth, development and maturity. It helps in flowering and fruiting. Phosphorus is a constituents of nucleic acid, phytine, phospholipids and majority of enzymes, which are of energy in carbohydrate and fat metabolism. It hastens maturity and improves quality of grain, therefore application of phosphorus is must. Potassium in cell sap is involved in enzyme activation, photosynthesis, transport of sugar, protein and starch synthesis. It is known to help crop to perform better under water stress condition through the regulation of the transpiration rate at which plant stomata open and close. It is also known for its role to provide lodging resistance, pest and disease resistance to plants.

Materials and Methods

The field experiments was conducted at Post Graduate Research farm Agronomy section, RSCM College of Agriculture, Kolhapur during March to April 2020 to study the effect of fertilizer levels and plant densities on growth attributes and yield of summer sunflower. The soil type was clay loam with pH of 7.75, organic matter 0.8%, Available nitrogen 236.08 kg ha⁻¹, phosphorus 27.50 kg ha⁻¹ and potassium 287.02 kg ha⁻¹. The experiment consisted nine treatments with a combination of three fertilizer levels (45:45:45, 60:60:60 and 75:75:75 kg NPK ha⁻¹) and plant densities (1,11,111; 74,074 and 55,555 plants ha⁻¹) in a Factorial randomized block design.

Crop was planted in a plot size of 25.92 m² (5.4m x 4.8m) at a spacing of 30 cm x 30 cm, 45 cm x 30 cm and 60 cm x 30 cm. The crop was planted in March 2020. The sunflower crop was fertilized with 45:45:45, 60:60:60 and 75:75:75 kg N: P₂O₅: K₂O ha⁻¹. Full dose of fertilizer were applied at the time of sowing. Nitrogen was applied through urea, phosphorus

through single super phosphate (SSP) and potassium through muriate of potash. Gap filling was done at 15th days after sowing and thinning was done at 24th days after sowing, two manual weeding were done throughout the sunflower growing period, irrigation was done four times during the growing period of sunflower.

Five plants were randomly selected from each plot for collecting data on growth attributes and yield. At full maturity, the crop was harvested plot-wise on 21st May and 31st May 2021. Seed and stalk was thoroughly dried plot by plot individually before their weights were recorded. Seed obtained from each unit plot were sun-dried and weighed carefully and the plot yield was recorded in quintal per hectare (q ha⁻¹). Stalk obtained from each unit plot were dried in sun and final stalk yield per plot was recorded in q ha⁻¹.

The experimental data was statistically analyzed by using a standard method of "analysis of variance" as reported by Panse and Sukhatme (1967) [17].

Result and Discussion

Growth attributes

The plant height⁻¹ (cm), number of functional leaves⁻¹, leaf area⁻¹ (dm²) and dry matter⁻¹(g), were significantly influenced by different fertilizer levels (Table 1). There was an increasing trend was observed in all the growth attributes with increasing fertilizer levels. Similar trend of growth attribute were reported by Maheshbabu *et al.*, (2007) [14]. Highest plant height (182.42 cm), number of functional leaves⁻¹ (22.42), leaf area⁻¹ (17.17 dm²) and dry matter⁻¹ (120.67 g) were obtained with 75:75:75 kg NPK ha⁻¹, however it was on par with the 60:60:60 kg NPK ha⁻¹ and lowest value were recorded with 45:45:45 kg NPK ha⁻¹ (Table 1). This might due to higher levels of NPK was result of enhanced activities of meristematic tissue of the plant, increase in number and size of the cell, and the efficient utilization of nutrient uptake. A similar results was also reported by Khokani *et al.*, (1992) [11] and Ali *et al.*, (2012) [11].

The planting density of 30 cm x 30 cm produced highest plant height (182.67 cm), however on par with the 45 cm x 30 cm and lowest plant height was recorded in 60 cm x 30 cm, which might be due to the competition among the plants get more sunlight. Similar results were also reported by Asghar Ali *et al.*, (2007) [2] and planting density 60 cm x 30 cm produced highest number of functional leaves⁻¹ (22.33), leaf area⁻¹ (16.42 dm²) and dry matter plant⁻¹ (120.33 g), however it was on par with the 45 cm x 30 cm and lowest value was recorded 30 cm x 30 cm (Table 1). This might due to less availability of space for aeration, expansion, radiation and more competition for available nutrients and moisture. Similar results reported by Bhoite and Nimbalkar (1955) [5] and Shedgi *et al.*, (2008).

Table 1: Effect of fertilizer levels and plant densities on growth attributes of summer sunflower

At harvest				
Treatments	Plant height ⁻¹ (cm)	Number of functional leaves ⁻¹	Leaf area ⁻¹ (dm ²)	Dry matter plant ⁻¹ (g)
Fertilizer levels (N: P ₂ O ₅ : K ₂ O Kg ha ⁻¹) (F)				
F1- 75% RDF (45:45:45)	168.58	19.33	10.25	109.67
F ₂ - 100% RDF (60:60:60)	177.08	20.75	16.00	118.42
F ₃ - 125% RDF (75:75:75)	182.42	22.42	17.17	120.67
S.E. ±	3.57	0.65	0.46	2.46
CD at 5%	10.43	1.92	1.37	7.20
Plant Density (D)				
D ₁ - 1,11,111 plants ha ⁻¹ (30 cm x 30 cm)	182.67	19.17	11.92	111.33

D ₂ – 74,074 plants ha ⁻¹ (45 cm x 30 cm)	176.75	21.00	15.08	117.08
D ₃ – 55,555 plants ha ⁻¹ (60 cm x 30 cm)	168.67	22.33	16.42	120.33
S.E. ±	3.57	0.65	0.46	2.46
CD at 5%	10.43	1.92	1.37	7.20
Interaction Effect (F x D)				
S.E. ±	6.19	1.14	0.81	4.27
CD at 5%	NS	NS	NS	NS
Grand Mean	176.02	20.84	14.47	116.25

Seed yield, stalk yield and Biological yield

Seed yield, stalk yield and biological yield were significantly influenced by different fertilizer levels (Table 2). The highest seed yield (13.88 q ha⁻¹), stalk yield (26.50 q ha⁻¹) and biological yield (40.38 q ha⁻¹) was obtained with 75:75:75 kg NPK ha⁻¹, however it was on par with the 60:60:60 kg NPK ha⁻¹ and lowest seed yield (11.25 q ha⁻¹), stalk yield (21.00 q ha⁻¹) and biological yield (32.25 q ha⁻¹) was achieved with 45:45:45 kg NPK ha⁻¹. The highest seed yield, stalk yield and biological yield was observed in application if 75:75:75 kg NPK ha⁻¹ it might due to more availability of plants nutrients leads to the healthy and vigorous growth of the crop. A similar results were also reported by Nasim *et al.*, (2011)^[15], Volkan and Kamaltin (2015)^[19] and Oad *et al.*, (2018)^[18].

Seed yield, stalk yield and biological yield were significantly influenced by different plant densities (Table 2). The seed

yield at planting density 60 cm x 30 cm was the highest (13.83 q ha⁻¹), however it was on par with the planting density 45 cm x 30 cm and the lowest one (11.33 q ha⁻¹) was recorded at 30 cm x 30 cm spacing (Table 2). Due to closer planting density seed yield gradually decreased. Under closer planting density, the rate of yield reduction was in response to decreasing radiation, nutrients, moisture and air. This result is in agreement with that of Kandil *et al.*, (2017)^[18]. The highest Stalk yield (28.42 q ha⁻¹) and biological yield (39.75 q ha⁻¹) was obtained with the 30 cm x 30 cm, however it was on par with the 45 cm x 30 cm and the lowest stalk yield (19.25 q ha⁻¹) and biological yield (33.08 q ha⁻¹) was recorded in the 60 cm x 30 cm, because least number of plants per unit area than the rest of planting densities. Similar results were observed by Gholinezhed *et al.*, (2009).

Table 2: Effect of fertilizer levels and plant densities on yield of summer sunflower

Treatments	Seed yield (q ha ⁻¹)	Stalk yield (q ha ⁻¹)	Biological yield (q ha ⁻¹)
Fertilizer Levels (N: P₂O₅: K₂O Kg ha⁻¹) (F)			
F ₁ - 75% RDF (45:45:45)	11.25	21.00	32.25
F ₂ - 100% RDF (60:60:60)	12.63	23.42	36.04
F ₃ - 125% RDF (75:75:75)	13.88	26.50	40.38
S.E. ±	0.50	1.00	1.14
CD at 5%	1.47	2.93	3.34
Plant Densities (D)			
D ₁ - 1,11,111 plants ha ⁻¹ (30 cm x 30 cm)	11.33	28.42	39.75
D ₂ – 74,074 plants ha ⁻¹ (45 cm x 30 cm)	12.58	23.25	35.83
D ₃ – 55,555 plants ha ⁻¹ (60 cm x 30 cm)	13.83	19.25	33.08
S.E. ±	0.50	1.00	1.14
CD at 5%	1.47	2.93	3.34
Interaction Effect (F x D)			
S.E. ±	0.87	1.74	1.98
CD at 5%	NS	NS	NS
Grand Mean	12.58	23.63	36.22

Conclusion

Based on the present study, it may be concluded that the application of 75:75:75 kg NPK ha⁻¹ found to be beneficial in increasing growth attribute and yield of summer sunflower. Sowing 60 cm x 30 cm spacing (planting density 55,555 plants ha⁻¹) recorded higher growth attribute and yield of summer sunflower.

Acknowledgement

The authors are thankful to Agronomy Section, RSCM College of Agriculture, Kolhapur – 416004, Maharashtra, India for providing necessary facilities to undertake the field experiment.

References

1. Ali A, Ahmad A, Khaliq T, Akhtar J. Planting density and nitrogen rates optimization for growth and yield of sunflower (*Helianthus annuus* L.) hybrids. The Journal of Animal and Plant Sciences 2012;22(4):1070-1075.
2. Ali A, Tanveer A, Nadeem MA, Tahir M, Hussain M.

Effect of varying planting pattern on growth, achene yield and oil contents of sunflower (*Helianthus annuus* L.). Pakistan Journal of Agricultural Sciences 2007;44:449-452.

3. Anonymous. Directorate of Economics and Statistics, Department of Agriculture and Cooperation, Ministry of Agriculture, GOI, 2019.
4. Basha HA. Response of two sunflower cultivars to hill spacings and nitrogen fertilizer levels under sandy soil conditions. Zagazig Journal of Agricultural Research (Egypt) 2000;27(4):617-633.
5. Bhoite SV, Nimbalkar VS. Effect of plant densities and fertilizer levels on yield of *kharif* sunflower. J Maharashtra Agric. Univ 1995;20(1):153-154.
6. Biabani A. Cultivar and density effects on yield of soybean in double cropping. African Journal of Agricultural Research 2010;5(23):3203-3206.
7. Gholinezhad E, Aynaband A, Ghorthapeh AH, Noormohamadi G, Bernousi I. Study of the effect of drought stress on yield, yield components and harvest

index of sunflower hybrid iroflor at different levels of nitrogen and plant population. Notulae Botanicae Horti Agrobotanici Cluj-Napoca 2009;37(2):85-94.

8. Kandil AA, Sharief AE, Odam AMA. Response of some sunflower hybrids (*Helianthus annuus* L.) to different nitrogen fertilizer rates and plant densities. International Journal of Environment, Agriculture and Biotechnology 2017;2(6):238990.
9. Khan AZ, Shah P, Khalil SK, Taj FH. Influence of planting date and plant density on morphological traits of determinate and indeterminate soybean cultivars under temperate environment. Pakistan Journal of Agronomy 2003;2(3):146-152.
10. Hossam Ibrahim M. Response of some sunflower hybrids to different levels of plant density. ICAAA: July 23-24, 2012, Singapore.
11. Khokani MG, Ahlawat PS, Trivedi SJ. Effect of nitrogen and phosphorus on growth and yield of sunflower (*Helianthus annuus* L.). Indian Journal Agron 1992;38(3):507-509.
12. Liu X, Jin J, Wang G, Herbert SJ. Soybean yield physiology and development of high-yielding practices in Northeast China. Field crops research 2008;105(3):157-171.
13. Long M, Feil B, Diepenbrock W. Effects of plant density, row spacing and row orientation on yield and achene quality in rainfed sunflower. Acta agronomica hungarica 2001;49(4):397-407.
14. Maheshbabu HM, Hunje R, Patil NB, Babalad HB. Effect of organic manures on plant growth, seed yield and quality of soybean. Karnataka Journal of Agricultural Sciences 2008;21(2):219-221.
15. Nasim W, Ahmad A, Wajid A, Akhtar J, Muhammad D. Nitrogen effects on growth and development of sunflower hybrids under agro-climatic conditions of Multan. Pak. J Bot 2011;43(4):2083-2092.
16. Oad RK, Ansari MA, Kumar J, Menghwar DR. Effect of foliar applied urea on growth and yield of sunflower (*Helianthus annuus* L.). Open Access Library journal 2018;5(7):1-13.
17. Panse VG, Sukhamate PV. Statistical method for agriculture workers, ICAR, New Delhi 1967, 361.
18. Sedghi M, Seyed Sharifi R, Namvar A, Khandan-e-Bejandi T, Molaei P. Responses of sunflower yield and grain filling period to plant density and weed interference. Research Journal of Biological Sciences 2008;3(9):1048-1053.
19. Volkan GUL, Kemalettin KARA. Effects of different nitrogen doses on yield and quality traits of common sunflower (*Helianthus annuus* L.). Turkish Journal of Field Crops 2015;20(2):159-165.