



ISSN (E): 2277- 7695

ISSN (P): 2349-8242

NAAS Rating: 5.23

TPI 2021; 10(4): 01-03

© 2021 TPI

[www.thepharmajournal.com](http://www.thepharmajournal.com)

Received: 03-02-2021

Accepted: 12-03-2021

**Ravshankhon Makhmudov**  
Drug Development Department,  
Nika Pharm Co., Tashkent,  
Republic of Uzbekistan

**Furkat Yunusov**  
Tashkent State Technical  
University, State Unitary  
Enterprise Science and Progress,  
Tashkent, Uzbekistan

## Simple and rapid RP-HPLC determination of nimesulide in tablets

**Ravshankhon Makhmudov and Furkat Yunusov**

DOI: <https://doi.org/10.22271/tpi.2021.v10.i4a.6437>


### Abstract

The present investigation describes a fast and simplified RP-HPLC method for the evaluation of nimesulide in tablets. The method is specific, accurate, precise, and ensures good reproducibility. The sample was analyzed using Acetonitrile (ACN): Ethanol: Water in the ratio of 40:40:20 as a mobile phase at a flow rate of 1.0 ml/min and detection at 276 nm. The retention time for nimesulide was found to be about 3.4, and a recovery from tablet was 99%. The method can be used for the fast and simple quantitative estimation of nimesulide in tablets.

**Keywords:** Chromatography, HPLC, Nimesulide, Dysmenorrhea

### 1. Introduction

Nimesulide is a non-selective non-steroidal anti-inflammatory drug (NSAID). It is used to treat acute (short-term) pain, painful osteoarthritis (swelling in the joints), primary dysmenorrhea (period pains). Its approved indications are the treatment of acute pain, the symptomatic treatment of osteoarthritis, and primary dysmenorrhea in adolescents and adults above 12 years old [1-3]. The IUPAC name of nimesulide is N-(4-nitro-2-phenoxyphenyl)-methanesulphonamide (Fig. 1).



**Fig 1:** Chemical structure of nimesulide

There have been published several articles to estimate the quantity of nimesulide using HPLC [4-6], spectrophotometric [7, 8] and IR methods [9], as well as dissolution profile, validation of analytical methods for capsules having nimesulide [10].

Our goal of present investigation was to find a proper and environmentally safe eluent as well as to decrease the retention time of nimesulide peak in HPLC.

### 2. Instrumentation

The modular Shimadzu Nexera2 HPLC system is comprised of the following parts, each of those is produced by Shimadzu USA Manufacturing, Inc.: an SIL-30AC autosampler, an LC-30AD liquid chromatograph, a CTO-30A column oven; a binary pump; a SPD-20AV prominence UV-Vis detector; a DGU-20A5R degassing unit and a CMB-20A communications bus module. The chromatographic parameters (peak areas, retention times, theoretical plates etc.) were recorded and calculated via Shimadzu LabSolutions v.5.73 software running under MS Windows 7. The HPLC experiments were performed on a stainless-steel column 150 mm long, 4.6 mm internal diameter filled with Octadecyl silane chemically bonded to porous silica particles (Inert Sustain C18, 5 $\mu$ , 250 mm x 4.6 mm, Japan). The chromatographic conditions are listed in Table 1 below.

**Corresponding Author:**  
**Ravshankhon Makhmudov**  
Drug Development Department,  
Nika Pharm Co., Tashkent,  
Republic of Uzbekistan

**Table 1:** Chromatographic conditions

Parameters	Chromatographic conditions
Chromatograph	Shimadzu Nexera2 HPLC
Column	InertSustain C18, 5 $\mu$ , 250 mm x 4.6 mm, Japan
Mobile phase	ACN/Ethanol/Water
Flow rate	1.0 ml/min

### 3. Experimental

#### 3.1 Chemicals and Materials

A pure sample of nimesulide was purchased from Aarti Drugs Limited (India). The HPLC grade orthophosphoric acid, acetonitrile and ethanol purchased from VWR Chemical (France). Water was purified by a Milli-Q water purification system (Millipore, Bedford, MA, USA).

#### 3.2 Preparation of standard stock solution

The standard stock solution of 2 mg/ml of nimesulide was prepared by dissolving 100 mg of the working standard of nimesulide in 35 ml of 96% ethanol, treated by an ultrasonic for 15 min, then the solution was adjusted to the volume of 50 ml with 96% ethanol.

The top of this paragraph illustrates a sub-subheading.

#### 3.3 Preparation of test solution

378 mg of powder prepared of 20 tablets containing 100 mg of nimesulide was accurately weighed and placed into a 50 ml volumetric flask, then 35 ml of 96% ethanol was added, treated by ultrasonic for 15 min, the solution volume was adjusted to 50 ml with 96% ethanol, to get a concentration of 2 mg/ml of nimesulide.

Prior to injection all solutions were filtered through 0.45  $\mu$  membrane filters (Whatman). 3  $\mu$ l of the solution was injected into HPLC system to obtain the chromatograms for the standard drug solution (five replicates) and the sample solution (five replicates). The concentrations of nimesulide in the formulation were calculated by comparing the AUC of the sample with that of the standard.

#### 3.4 HPLC Procedure

All solutions of samples and standard (3 $\mu$ l) were injected by means of an autosampler and detected at 276 nm. ACN/Ethanol/Water (40/40/20) was used as the mobile phase while the flow rate was set at 1.0 ml/min at room temperature. Under the above-mentioned conditions, the analysis cycle was completed in 5 min.


## 4. Results

### 4.1 Validation of the HPLC Method

The fast and precise estimation of nimesulide in tablets by HPLC method was validated for linearity, precision (repeatability and reproducibility), and accuracy.

### 4.2 Linearity

The linearity and range of the method was determined by analyzing five different concentrations of standard solution ranging from 1.6 – 2.4 mg/ml. The calibration curve was plotted using the peak area versus the concentration of standard solution and correlation coefficient and regression line equation was determined. The linear plot of nimesulide is given in Fig. 2.



**Fig 2:** Calibration curve of nimesulide at 276 nm

### 4.3 Accuracy

Accuracy values shows that the estimated values are close to the reference ones prepared in laboratory, since in the range of 80% - 120%, the accuracy of the HPLC method for the quantitative determination of nimesulide was confirmed by appropriate tests of model solutions, based on the calculated results of the recovery rate. The results obtained are within the acceptable range of 98.0% - 102.0%. The average value of the degree of recovery was 100.16% of nimesulide (Table 2). Therefore, it is possible to conclude that these is acceptable for quality control of nimesulide in pharmaceutical formulation.

**Table 2:** The average value of the degree of recovery was 100.16% of nimesulide

Active compound	№	Quantity entered [mg] and [% of theoretical stated]		Detected amount, mg	Recovery rate, %
		Quantity [mg]	Quantity [%]		
Nimesulide	1	80	80%	79.76	99.70
	2	80	80%	80.17	100.21
	3	80	80%	80.03	100.04
	4	100	100%	100.24	100.24
	5	100	100%	100.53	100.53
	6	100	100%	100.40	100.40
	7	120	120%	120.34	100.28
	8	120	120%	119.83	99.86
	9	120	120%	120.23	100.19
Average recovery rate [%]					100.16
95% confidence interval					100,16 $\pm$ 0,16
Standard deviation					0.26
RSD [%]					0.26


#### 4.4 Convergence

The convergence of the method was studied twice in two days by using six determinations at 100% test concentration, i.e. mixture of 2 mg/ml of nimesulide. The results are given in Table 3.

**Table 3:** The results of convergence study

Active compound	S. №	The quantitative content of the model drug from 100% of theoretical one	
		1st day	2nd day
Nimesulide	1	100.66	100.88
	2	100.74	100.32
	3	100.44	101.46
	4	99.74	101.18
	5	99.96	101.67
	6	99.83	101.38
Average		100.23	101.15
Standard deviation		0.44	0.49
RSD [%]		0.438	0.481
95% confidence interval		100,23 ±0,357	101,15 ±0,39

The obtained test results of nimesulide, performed by six determinations on different days, reveals that the relative standard deviations of the RSD are 0.438% and 0.481%, which indicates the acceptable degree of convergence of the quantitative content of model preparations (Fig 3).



**Fig 3:** Chromatogram of nimesulide

#### 5. Conclusions

The following validation characteristics have been confirmed for the validation of the quantitative determination method for nimesulide:

1. Specificity
2. Accuracy
3. Linearity
4. Convergence

Their compliance with the established acceptance criteria was experimentally proved.

These analytical methods should be considered validated, they correspond to the intended purpose: to determine the authenticity and quantitative determination of nimesulide. They can be used for quality control of this medicinal product.

#### 6. References

1. Rainsford KD. Nimesulide - Actions and Uses;

Birkhauser Verlag, Bassel 2005.

2. Goodman LS, Gilman A. The Pharmacological Basis of Therapeutics, 9th ed., Macmillan Publishing Co. New York 1975.
3. Rowe RC, Sheskey PJ, Owen SC. Handbook of Pharmaceutical Excipients, 5th Ed., Pharmaceutical Press, London; Chicago. Washington DC 2006.
4. Prasanna Reddy Battu. Determination of Nimesulide in Pharmaceutical formulations and in Human serum by Reverse-Phase High-Performance Liquid Chromatography Int. J Pharm Tech Res 2009;1(3)
5. Zarapkar SS, Bhandari NP, Halkar UP. Simultaneous determination of nimesulide and chlorzoxazone in pharmaceutical dosage by RP-HPLC. Indian Drugs 2000;37:467.
6. Nagoji KE, Vijaysrinivas S, Kumar KM, Mathivanan N, Kumar SM, Rao ME. Simultaneous RP-HPLC estimation of nimesulide and diclofenac sodium. Indian J Pharma Sci 2003;65:407.
7. Rajput SJ, Randive G. Assay of nimesulide in pharmaceutical dosage formulation. Eastern Pharmacist 1997;475:113.
8. Nagoji KE, Rao SS, Rao ME, Rao KV. New Spectroscopic method for the estimation of Nimesulide in Pharmaceutical Dosage formulation. Eastern Pharmacist 1999;496:117.
9. Werickson FC Rocha, André Luis Rosa, José Antônio Martins, Ronei Jesus Poppi. Determination and validation of nimesulide in pharmaceutical formulation by near infrared spectroscopy. J Braz. Chem Soc 2010;21(10):1929-1936.
10. Beatriz Werneck Lopes Santos, Eloisa Dutra Caldas, Mônica Valero Da Silva. Nimesulide: Dissolution Profile, Validation of Analytical Methods for Capsules, and Assessment of Product Quality. Int. J Pharm Pharm Sci 2016;8(1):27-32.