



ISSN (E): 2277- 7695  
ISSN (P): 2349-8242  
NAAS Rating: 5.23  
TPI 2021; 10(6): 1015-1017  
© 2021 TPI

[www.thepharmajournal.com](http://www.thepharmajournal.com)

Received: 14-03-2021

Accepted: 23-05-2021

## Neeraj Singh

Department of Agriculture  
Statistics, Acharya Narendra  
Deva University of Agriculture  
& Technology, Kumarganj,  
Ayodhya, Uttar Pradesh, India

## Piyush Kumar Singh

Department of Agriculture  
Statistics, Acharya Narendra  
Deva University of Agriculture  
& Technology, Kumarganj,  
Ayodhya, Uttar Pradesh, India

## VN Rai

Department of Agriculture  
Statistics, Acharya Narendra  
Deva University of Agriculture  
& Technology, Kumarganj,  
Ayodhya, Uttar Pradesh, India

## BVS Sisodia

Department of Agriculture  
Statistics, Acharya Narendra  
Deva University of Agriculture  
& Technology, Kumarganj,  
Ayodhya, Uttar Pradesh, India

## Corresponding Author:

### Neeraj Singh

Department of Agriculture  
Statistics, Acharya Narendra  
Deva University of Agriculture  
& Technology, Kumarganj,  
Ayodhya, Uttar Pradesh, India

## Effect of weather variables on the yield of wheat crop in District Azamgarh of Eastern Uttar Pradesh, India

Neeraj Singh, Piyush Kumar Singh, VN Rai and BVS Sisodia

### Abstract

Wheat (*Triticum aestivum* L.) is the major crop of Uttar Pradesh, which covers about 36.5 per cent area of total gross-cropped area in Uttar Pradesh. The present study mainly deals with to determine the effect of weather variables on the yield of Wheat crop in Azamgarh district of Eastern Uttar Pradesh, India. On the basis of correlation and regression analysis, we found that rainfall is most important variable which effects the yield of Wheat crop positively followed by minimum temperature while rest variables show negative effects on the yield of Wheat crop.

**Keywords:** Azamgarh, wheat, yield, weather variables, correlation, regression, r-square

### Introduction

Wheat (*Triticum aestivum* L.) is the major crop of Uttar Pradesh, which covers about 32.89 per cent area of total gross-cropped area and contributes 31.92 per cent production of India in 2017-18, (DAC&FW, 2018). Wheat is mostly grown in Rabi (December-April) season. This crop is also studied under the FASAL programme of the Department of Space (Parihar & Oza, 2006). Due to importance of this crop in the national agricultural scenario, it is important to study the impact of climate change on this crop.

Azamgarh district (Eastern Uttar Pradesh, India) which falls under middle gangetic agro-climatic region and 8<sup>th</sup> eastern plain agro-climatic zone in the eastern part of Uttar Pradesh at a distance of about 270 km from Lucknow. Azamgarh district lies between latitude 26° 03' N and longitude 83° 13' E and it is bounded by the districts of Mau in east, Gorakhpur in the north, Ghazipur in south-east, Jaunpur in the south-west, Sultanpur in the west and Ambedkar Nagar in the north-west.

The present study has been undertaken for the yield of Wheat crop in the district Azamgarh, Eastern Uttar Pradesh, India.

### Material and Methods

The time series data on yield for Wheat crop of Azamgarh district of eastern Uttar Pradesh pertaining for the period from 2000-01 to 2017-18 have been procured from the website <http://updes.up.nic.in/spatrika/spatrika.htm> by Economics and Statistics Division, Planning Department, Government of Uttar Pradesh.

Weekly weather variables data for Wheat and wheat crop in the district of Azamgarh, Eastern Uttar Pradesh have been obtained from the National Data Centre, India Meteorological Department, Pune for the study period 2000-01 to 2017-18. The data for Wheat crop have been collected up to the first 16 weeks of the crop cultivation which include 44<sup>th</sup> Standard Meteorological Week (SMW) to 7<sup>th</sup> SMW of the next year. The data on six weather variables viz. Maximum Temperature, Minimum Temperature, Rainfall, wind-velocity and Sun-shine hours have been used in the study.

### Individual effect of weather variables

The statistical models have been proposed by expressing effect of changes in weather variables on yield in w<sup>th</sup> week as a linear function of respective correlation coefficients between detrended yield and weekly weather data (Agrawal *et al.*, 1986). Trend effect on yield is also removed from yield while calculating correlation coefficients of yield with weather variables to be used as weights.

**De-trend Yield**

$$Y = a + bt$$

Where; Y, a, b and t is observed yield, constant, regression coefficient and time trend respectively.

In order to study, the effect of individual weather variable, two new variables from each weather variable are generated as follows:

Let  $X_{iw}$  be the value of  $i^{th}$  ( $i = 1, 2, \dots, p$ ) weather variable at  $w^{th}$  weeks ( $w = 1, 2, \dots, n$ ). In this study, n is 16.

Let,  $r_{iw}$  be the simple correlation coefficient between weather variable  $X_i$  at W-th week and detrended crop yield over a period of K years. The generated variables are then given by

$$Z_{ij} = \frac{\sum_{w=1}^n r_{iw}^j X_{iw}}{\sum_{w=1}^n r_{iw}^j}; j = 0, 1$$

For  $j = 0$ , we have un-weighted generated variable

$$Z_{i0} = \frac{\sum_{w=1}^n X_{iw}}{n}$$

and weighted generated variables

$$Z_{i1} = \frac{\sum_{w=1}^n r_{iw} X_{iw}}{\sum_{w=1}^n r_{iw}}$$

For each year.

The following model is then fitted to study the effect of individual weather variable

$$Y = a_0 + a_1 Z_{i0} + a_2 Z_{i1} + cT + \epsilon; i = 1, 2, \dots, p.$$

Where, Y is detrended yield. T is variable expressing time effect,  $a_0, a_1, a_2$  and c are parameters of the model to be evaluated for the effect of variables and  $\epsilon$  is error term supposed to follow normal distribution with mean zero and variance  $\sigma^2$ .

**Result and Discussion**

**Effect of weather variables on the yield of Wheat crop by Correlation Analysis**

Persual of the table 1, we found that unweighted minimum temperature ( $Z_{20}$ ) and unweighted rainfall ( $Z_{30}$ ) are positively correlated as 0.560 and 0.531 respectively with detrended yield of Wheat crop at 5% level of significance while unweighted maximum temperature ( $Z_{10}$ ), unweighted wind velocity ( $Z_{40}$ ) and unweighted sunshine hour ( $Z_{50}$ ) found to be negatively correlated with detrended yield of Wheat crop.

**Table 1:** Correlation Coefficient between Detrend yield and generated Weather variables

Unweighted Variables	Correlation Coefficient	Weighted Variables	Correlation Coefficient
$Z_{10}$	- 0.259	$Z_{11}$	- 0.621**
$Z_{20}$	0.560*	$Z_{21}$	0.441
$Z_{30}$	0.531*	$Z_{31}$	0.738**
$Z_{40}$	- 0.398	$Z_{41}$	0.167
$Z_{50}$	- 0.390	$Z_{51}$	- 0.641**

Correlation is significant at the 0.01 level (2-tailed) \*\*

Correlation is significant at the 0.05 level (2-tailed) \*

Weighted minimum temperature ( $Z_{21}$ ), weighted rainfall ( $Z_{31}$ ) and weighted wind velocity ( $Z_{41}$ ) shows positive while

weighted maximum temperature ( $Z_{11}$ ) and sunshine hour ( $Z_{51}$ ) shows negative correlation with the yield of Wheat crop.

**Effect of weather variable on the yield of Wheat crop by Regression Analysis**

**Effect of Maximum Temperature**

The multiple regression equation obtained is

$$Y = 63.013 - 0.331 Z_{10} - 1.011 Z_{11} - 0.143 T$$

**Table 2:** Effect of Maximum Temperature

Variable	Regression Coefficient (Standard Error)	P Value	R <sup>2</sup> (%)	95% Confidence Interval	
				Lower B.	Upper B.
Constant	63.013*** (18.312)	0.004	58.6	23.738	102.287
$Z_{10}$	- 0.331 (0.509)	0.527		- 1.423	0.761
$Z_{11}$	- 1.011*** (0.238)	0.001		- 1.522	- 0.500
T	- 0.143 (0.101)	0.180		- 0.360	0.074

From the persual of table 2 maximum temperature is an important weather variable for Wheat crop. The results indicates that weighted maximum temperature regression coefficients were found to be statistically negatively significant at 1% level of significant and time trend T was found negatively non-significant. The value of  $R^2$  (%) is 58.6 and  $F_{Cal}$  (6.614\*\*\*) is significant at 5% level of significance.

**Effect of Minimum Temperature**

The multiple regression equation obtained is

$$Y = 21.685 - 0.181 Z_{20} + 0.205 Z_{21} - 0.045 T$$

**Table 3:** Effect of Minimum Temperature

Variable	Regression Coefficient (Standard Error)	P Value	R <sup>2</sup> (%)	95% Confidence Interval	
				Lower B.	Upper B.
Constant	21.685** (9.055)	0.031	12.3	2.264	41.106
$Z_{20}$	- 0.181 (0.848)	0.834		- 1.999	1.638
$Z_{21}$	0.205 (0.150)	0.193		- 0.116	0.525
T	- 0.045 (0.135)	0.745		- 0.334	0.244

From the persual of table 3 indicates that un-weighted & weighted minimum temperature was found statistically negative and positive non-significant respectively and time trend T was found negatively non-significant. The value of  $R^2$  (%) is 12.3,  $F_{Cal}$  (0652) is non-significant.

**Effect of Rainfall**

The multiple regression equation obtained is

$$Y = 23.995 - 0.793 Z_{30} - 0.195 Z_{31} - 0.127 T$$

**Table 4:** Effect of Rainfall

Variable	Regression Coefficient (Standard Error)	P Value	R <sup>2</sup> (%)	95% Confidence Interval	
				Lower B.	Upper B.
Constant	23.995 (1.408)	.000	48.8	20.975	27.016
$Z_{30}$	-0.793** (0.314)	0.024		- 1.468	- 0.119
$Z_{31}$	-0.195*** (0.059)	0.005		- 0.321	- 0.068
T	-0.127 (0.101)	0.230		- 0.343	0.090

From the persual of table 4, un-weighted & weighted rainfall regression coefficient was found statistically negative significant at 5% and 1% respectively, where time trend T was found non-significant. The value of  $R^2$  (%) is 48.8 and  $F_{Cal}$  (4.44\*\*) is significant.

## 2.4 Effect of Wind Velocity

The multiple regression equation obtained is

$$Y = 22.679 - 1.552 Z_{40} + 1.344 Z_{41} - 0.018 T$$

**Table 5:** Effect of Wind velocity

Variable	Regression Coefficient (Standard Error)	P Value	R <sup>2</sup> (%)	95% Confidence Interval	
				Lower B.	Upper B.
Constant	22.679 (4.333)	0.000	40.4	13.386	31.972
Z <sub>40</sub>	- 1.552 (1.050)	0.162		- 3.803	0.700
Z <sub>41</sub>	1.344** (0.459)	0.011		0.360	2.329
T	- 0.018 (0.118)	0.878		- 0.272	0.235

From the persual of table 5, Un-weighted regression coefficient of wind velocity was found to be negatively non-significant, weighted variable was found positively significant at 5% and time trend was negatively non-significantly. The value of R<sup>2</sup> (%) is 40.4 and F<sub>Cal</sub> (3.166\*) is significant at 10% level of significance.

## 2.5 Effect of Sunshine (hr)

The multiple regression equation obtained is

$$Y = 17.268 + 0.264 Z_{50} + 0.293 Z_{51} + 0.140 T$$

**Table 6:** Individual effect of Sunshine (hr)

Variable	Regression Coefficient (Standard Error)	P Value	R <sup>2</sup> (%)	95% Confidence Interval	
				Lower B.	Upper B.
Constant	17.268 (2.880)	.000	14.0	11.090	23.445
Z <sub>50</sub>	0.264 (0.503)	0.608		-.815	1.342
Z <sub>51</sub>	0.293 (0.322)	0.378		-.398	.984
T	0.140 (0.157)	0.386		-.196	.476

From the persual of table 6, weighted regression coefficient of sunshine hour was found negatively significant at 1% level, while regression coefficient of un-weighted sunshine hour and time was found negatively non-significant. The value of R<sup>2</sup> (%) is 14.0 and F<sub>Cal</sub> (0.759) is non-significant.

## Conclusion

All the weather variables which are used in this study is found to be important for the yield of Wheat crop in district Azamgarh, Eastern Uttar Pradesh, India. Correlation and regression analysis shows that among all these variables two variables *viz.*, rainfall and minimum temperature effects positively to the yield of the Wheat crop while all the rest variables effected negatively.

## Reference

1. Kumar N, Pisal RR, Shukla SP, Pandey KK. Crop yield forecasting of paddy, sugarcane and wheat through linear regression technique for south Gujarat. *Mausam* 2014;65(3):361-364.
2. Mehta SC, Agrawal R, Singh VPN. Strategies for composite forecast. *Jr Indian Soc. Ag. Stat* 2000;53(3):262-72.
3. Pandey KK, Kaushal RP, Mishra AN, Rai VN. A study of Impact of Weather Variables with Different Distributions. *Ind. J Agricult. Stat. Sci* 2009;5(1):139-53.
4. Pandey KK, Rai VN, Sisodia BVS. Weather Variables Based Wheat Yield Forecasting Models for Faizabad District of Eastern U.P. *Int. J Agricult. Stat. Sci*

2014;10(2):381-385.

5. Rai T, Chandrahas. Use of discriminant function of weather parameters for developing forecast model of Wheat crop. (IASRI Publication) 2000.
6. Ramasubramanian V, Jain RC. Use of growth indices in Markov chain model for crop yield forecasting. *Biometrical Journal* 1999;41(1):99-109.
7. Robertson GW. Wheat yields for 50 years at Swift Current Saskatchewan in relation to weather. *J Plant Sci* 1974;54:625-50.
8. Saksena A, Jain RC, Yadav RL. Development of early warning and yield assessment models for rainfed crops based on agro-meteorological indices. (IASRI Publication) 2001.
9. Shankar U, Gupta BRD. Forecasting the yield of paddy at chisurah in West Bengal using multiple regression technique. *Mausam* 1987;38:415-418.
10. Srivastava AK, Bajpai PK, Yadav RL, Hasan SS. Weather Based Sugarcane Yield Prediction Model for the State of Uttar Pradesh. *Journal of Indian Society of Agricultural Statistics* 2007;61(3):313-327.
11. Varshneya MC, Chinchorkar SS, Vaidya VB, Pandey V. Forecasting model for seasonal rainfall for different regions of Gujarat. *J Agrometeorology* 2010;12(2):202-207.