



ISSN (E): 2277- 7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2021; 10(7): 304-306
© 2021 TPI
www.thepharmajournal.com
Received: 07-04-2021
Accepted: 16-05-2021

SP Kushwaha
Department of Crop Physiology,
Chandra Shekhar Azad
University of Agriculture and
Technology, Kanpur, Uttar
Pradesh, India

DK Tripathi
Department of Crop Physiology,
Chandra Shekhar Azad
University of Agriculture and
Technology, Kanpur, Uttar
Pradesh, India

OP Singh
Department of Crop Physiology,
Chandra Shekhar Azad
University of Agriculture and
Technology, Kanpur, Uttar
Pradesh, India

Laxmi Prajapati
Department of Crop Physiology,
Chandra Shekhar Azad
University of Agriculture and
Technology, Kanpur, Uttar
Pradesh, India

Corresponding Author:
SP Kushwaha
Department of Crop Physiology,
Chandra Shekhar Azad
University of Agriculture and
Technology, Kanpur, Uttar
Pradesh, India

Effect of growth regulators on yield of paddy (*Oryza sativa L.*)

SP Kushwaha, DK Tripathi, OP Singh and Laxmi Prajapati

Abstract

The present investigation entitled “Effect of Growth Regulators on Yield of Paddy (*Oryza sativa L.*)” was conducted during the *kharif* season in 2012 and 2013 at Student’s Instructional Farm of Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P.). The experiment was setup in Randomized Block Design (RBD) with three replications and nine treatments. Growth regulators *i.e.* IAA (25 and 50 ppm), CCC (2000 and 4000 ppm), Alar (2000 and 3000 ppm) and Cytokinin (5 and 10 ppm) were applied on foliage at vegetative and milky stage of Rice (*Var.* PHB-71). Growth parameters were recorded at maximum maturity stage of paddy crop, while, Yield attributes were taken at the time of crop harvest. Chlorophyll Intensity (%) was recorded at pre-anthesis stage. All PGRs showed superior value in all the parameters of crop with respect to control. Growth characters *viz.* Plant Height (cm), Number of tillers plant⁻¹, Leaf area plant⁻¹ (cm²), Total dry matter production plant⁻¹ (g) were found superior with foliar application of IAA 50 ppm and Yield attributes *viz.* Number of panicles plant⁻¹, Number of grains panicle⁻¹, Grain weight plant⁻¹ (g), Grain yield (kg ha⁻¹), Test weight (g), Harvest index (%) were also maximized with foliar application of IAA 50 ppm. Chlorophyll intensity was found superior by the treatment of IAA 50 ppm.

Keywords: PGRs, paddy, growth, chlorophyll, yield

Introduction

Paddy (*Oryza Sativa L.*) is one of the most important cereal crops of the world in terms of food, area and production (Niamatullah *et al.* 2010) [6]. One in every three people depends on paddy for more than half of their daily food (Khush and Virk, 2000) [4]. The crop occupies one of the world’s total areas planted to cereals, providing 50-60 percent of the calories to 2.7 billion peoples (Belder *et al.* 2004) [2]. The demand of paddy is growing due to increasing population and food consumption patterns, while the land will either remain same or may decline. So, with the growing demand, it is very important to ensure the constant availability of more paddy at less cost. Therefore, the use of growth regulators not only provided the useful tool to improve the morphological but also helpful in overcoming the problem of lower yield of paddy crop. Plant growth regulators have been used in agriculture for as long as crop cultivation, their impact up to now has been relatively little detected and their application is limited to some specific objectives for example quality and quantity improvement (Pandey *et al.* 2001) [7]. PGRs play vital roles in coordination of many growth and behavioral processes in paddy, which regulates the amount, type and direction of plant growth (Rajendra and Jones Jonathan 2009; Anjum *et al.* 2011) [1, 8]. It modifies growth and development in various ways under different growth conditions and are effective in elevating the stress effects (Razi and Sen, 1996) [9]. It plays an important role in the movement of accumulated carbohydrate to the sink. Auxins (IAA) play a dominant role in root and shoot development, leaf senescence, fruit setting and flowering (Friml, 2003) [3]. Adjustment of plant growth and development using growth reducers such as Cycocel to reduce the plant height and increase the tolerance to lodging is a technique permitting the greater amount of nitrogen to attain higher yield (Mohaghegh *et al.* 2007) [5]. Cycocel treatment decreases the cell size and increases the cell wall thickness, thickness of the cell sap, increases the number of stem sets, decreases the length of internodes and increases the stem diameter. It decreases the plant height up to 23% and significantly increases the grain yield (Singh *et al.* 2002) [10]. Alar is a growth retardant used to reduce the plant height, increase stem diameter, short foliage stature of the crop plant. Cytokinin is used to promote cell division in plant root and shoot, improve plant height, chlorophyll content and grain filling rate.

Materials and Methods

The experiment was conducted at Student's Instructional Farm of Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P.). Geographically, Kanpur is located at 26.30° N Longitude, 80.15° E Latitude and it lies 127 meters above mean sea level in Gangatic alluvial of central Uttar Pradesh. The experimental site having sandy-loam soil with pH 7.2, Organic matter 0.46%, available nitrogen 245 kg/ha, available phosphorous 16 kg/ha and available potassium 282 kg/ha. The experiment was laid down in Randomized Block Design (RBD) with three replications. Paddy variety PHB-71 was sown in puddled bed to raise the nursery. Transplanting of twenty one days old seedlings with two seedlings per hill was done with 20 x 15 cm spacing (Plot size was 3 x 4 m²) in each three replications. Nitrogen, Phosphorus and Potash were applied in transplanted field at the rate of 120:60:60 kg ha⁻¹ in the form of urea, single super phosphate and murate of potash, respectively. Various plant growth regulators *viz.* IAA (25 and 50 ppm), CCC (2000 and 4000 ppm), Alar (2000 and 3000 ppm) and Cytokinin (5 and 10 ppm) were used as foliar application at vegetative and milky stage.

Observations on Growth characters *viz.* Plant Height (cm), Number of tillers plant⁻¹, Leaf area plant⁻¹ (cm²), Total dry matter production plant⁻¹ (g) were recorded at maturity stage of crop and Chlorophyll intensity (%) was measured with the help of Plant Efficiency Analyzer (Model X 55/M-PEA) and the values were expressed as SPAD value and it was measured at pre-anthesis stage. Yield with its attributes *viz.* Number of panicles plant⁻¹, Number of grains panicle⁻¹, Grain weight plant⁻¹ (g), Grain yield (kg ha⁻¹), Test weight (g), Harvest index (%) were taken at the time of harvesting.

Results and Discussion

A perusal of data on growth parameters presented in Table-1 revealed that application of growth regulators as foliar spray significantly influenced the plant height. The effect of IAA 50 ppm was more pronounced followed by IAA 25 ppm and Cytokinin 5 and 10 ppm in both years. While minimum plant height was recorded with treatment of CCC 4000 ppm. Both the concentration of CCC and Alar reduces the plant height in comparison to control. Data regarding Number of tillers plant⁻¹ (Table-1) as affected by growth regulators recorded at maturity stage. It is evident from the data that all the treatments significantly increased number of tillers Plant⁻¹.

The maximum Number of tillers plant⁻¹ was recorded with IAA 50 ppm followed by CCC 2000 ppm and IAA 25 ppm during 2012 and 2013 while minimum number of tillers plant⁻¹ was recorded with Cytokinin 10 ppm after controlled plant. Data pertaining to total dry matter production plant⁻¹ (g) (Table-1) as affected by foliar application of plant growth regulators cleared that the foliar application of all the treatments significantly increased total dry matter production plant⁻¹ (g) as compared to control during both years. The effect of foliar application of IAA 50 ppm was more pronounced and significantly increased the total dry matter production plant⁻¹ as compared to other treatments while minimum value was recorded with CCC 2000 ppm after control during 2012 and 2013. Result on leaf chlorophyll Intensity present in Table-1 represented that the growth regulators significantly accelerate the chlorophyll content of paddy leaf in comparison to control plant. The highest value on leaf chlorophyll was registered by the treatment of IAA 50 ppm followed by Cytokinin 10 ppm. Increment in total chlorophyll content was due to formation, maintenance and development of chloroplast.

Observation with respect to yield attributing characters showed that IAA 50 ppm registered significantly higher number of panicles plant⁻¹ (Table-2) as compared to other treatments including control during both years. Perusal of data revealed that the maximum number of grains panicle⁻¹ was recorded with IAA 50ppm followed by IAA 25 ppm and Alar 2000 ppm in 2011. Similar pattern was noted during 2012 and 2013. Data pertaining to grain weight plant⁻¹ (Table-2) as affected by foliar application of plant growth regulators. The maximum grain weight plant⁻¹ was recorded with IAA 50 ppm followed by Alar 2000 ppm and IAA 25 ppm. The data with respect to test weight (Table-3) revealed that the maximum test weight was recorded with IAA 50 ppm which found superior over rest of the treatments including control. The effect of foliar application of IAA 50 ppm was more pronounced and increased the grain yield (kg ha⁻¹) (Table-3) as compared to other treatments while minimum value was recorded under controlled plant during 2012 and 2013 of the research work. It is evident from the data that maximum harvest index (Table-3) was recorded with foliar application of IAA 50 ppm during 2011 while in 2012 it was maximum with Alar 2000 ppm and in 2013 maximum harvest index was found with Alar 3000 ppm.

Table 1: Effect of growth regulators on plant height, number of tillers plant⁻¹, total dry matter production plant⁻¹ and chlorophyll intensity (%) of paddy at maturity stage during 2012 and 2013

Treatment	Plant height (cm)		Number of tillers plant-1		TDM plant-1 (g)		Chlorophyll intensity (%)	
	2012	2013	2012	2013	2012	2013	2012	2013
T1 - Control	82.00	81.00	6.80	6.50	37.35	36.40	36.20	35.85
T2 - IAA 25 ppm	86.40	85.00	9.20	9.20	52.20	50.10	37.85	37.55
T3 - IAA 50 ppm	88.50	87.40	10.00	9.90	55.10	54.20	41.20	40.85
T4 - CCC 2000 ppm	77.70	76.60	9.50	9.10	42.50	43.00	36.75	35.95
T5 - CCC 4000 ppm	75.40	70.00	8.60	8.10	44.80	44.10	37.10	36.20
T6 - Alar 2000 ppm	80.20	79.40	9.00	8.20	51.60	48.20	36.85	36.35
T7 - Alar 3000 ppm	78.30	78.10	8.60	8.20	49.20	46.00	37.10	36.80
T8 - Cytokinin 5 ppm	85.10	83.60	8.90	8.50	50.80	48.30	37.25	37.00
T9 - Cytokinin 10 ppm	82.10	81.40	8.00	7.76	48.60	45.00	39.65	37.95
SE (m)±	1.13	1.29	0.29	0.73	1.02	1.39	2.03	1.88
CD at 5%	2.40	2.73	0.60	1.56	2.17	2.94	4.30	3.99

Table 2: Effect of growth regulators on number of panicles plant⁻¹, number of grains panicle⁻¹ and grain weight plant⁻¹ of paddy at harvesting stage during 2012 and 2013

Treatment	Number of panicles plant ⁻¹		Number of grains panicle ⁻¹		Grain weight plant ⁻¹ (g)	
	2012	2013	2012	2013	2012	2013
T1 - Control	6.80	6.50	134.33	131.33	20.00	19.70
T2 - IAA 25 ppm	9.00	8.60	160.22	158.00	23.00	23.00
T3 - IAA 50 ppm	10.00	9.00	170.20	165.00	25.40	25.10
T4 - CCC 2000 ppm	8.90	8.80	140.33	140.33	21.15	21.00
T5 - CCC 4000 ppm	8.00	8.00	135.66	136.66	20.10	20.00
T6 - Alar 2000 ppm	8.67	8.60	156.33	152.33	23.20	23.10
T7 - Alar 3000 ppm	8.33	8.20	152.66	150.66	22.00	22.00
T8 - Cytokinin 5 ppm	8.45	8.30	150.66	151.33	22.40	22.30
T9 - Cytokinin 10 ppm	8.08	7.80	147.70	148.66	21.60	21.40
SE (m)±	0.27	0.21	1.15	3.29	0.74	0.21
CD at 5%	0.55	0.43	2.43	6.97	1.56	0.45

Table 3: Effect of growth regulators on test weight, grain yield, harvest index of paddy at harvesting stage during 2012 and 2013

Treatment	Test weight (g)		Grain yield (kg ha ⁻¹)		Harvest index (%)	
	2012	2013	2012	2013	2012	2013
T1 - Control	20.00	19.40	3940.00	3848.00	31.43	30.00
T2 - IAA 25ppm	22.20	22.10	4276.00	4132.00	32.89	31.78
T3 - IAA 50ppm	23.20	23.20	4516.00	4420.00	34.74	33.23
T4 - CCC 2000ppm	20.40	20.10	4228.00	4113.00	32.11	31.31
T5 - CCC 4000ppm	20.25	20.00	4164.00	3992.00	31.03	30.84
T6 - Alar 2000ppm	22.65	22.40	4292.00	4180.00	32.52	31.66
T7 - Alar 3000ppm	21.55	21.60	4036.00	4084.00	31.20	31.17
T8 - Cytokinin 5ppm	21.45	21.35	4215.00	4132.00	33.63	32.53
T9 - Cytokinin 10ppm	20.00	20.45	4189.00	3940.00	33.51	31.35
SE (m)±	0.68	0.30	46.00	74.00	1.16	1.02
CD at 5%	1.44	0.64	99.00	152.00	2.46	2.20

References

1. Anjum SA, Wang LC, Farooq M, Hussain M, Xue LL, Zou CM. Brassinolide application improves the drought tolerance in maize through modulation of enzymatic antioxidants and leaf gas exchange. *J Agron. Crop Sci* 2011;197:177-185.
2. Belder P, Bouman BAM, Cabangon R, Guoan L, Quilang EJP, Yuanhua L *et al.* Effect of water saving irrigation on paddy yield and water use in typical low land conditions in Asia. *Agric. Water Manage* 2004;65(3):193-210.
3. Friml J. Auxin transport - shaping the plant. *Curr. Opin. Plant Biol* 2003;6(1):7-12.
4. Khush GS, Virk PS. Paddy breeding achievements and future strategies. *Crop Improvement* 2000;27(2):115-144.
5. Mohaghegh R, Imam Y. Study on the effect of Cycocel hormone on two cultivars of fall rapeseed. Thesis for M.S. in Shiraz University 2007.
6. Niamatullah M, Zaman KU, Khan MA. Impact of support paddy and fertilizer off take n paddy production and wheat acreage in NWFP, Pakistan. *J Animal Plant Sci* 2010;20(1):28-33.
7. Pandey AK, Tripathi RS, Yadav RS. Effect of certain growth regulators on growth yield and quality of paddy (*Oryza Sativa* L.). *Indian J Agric. Res* 2001;35(2):118-120.
8. Rajendra B, Jones Jonathan DG. Role of plant hormones in plant defense responses. *Plant Mol. Biol* 2009;69:473-488.
9. Razi SS, Sen SP. Diazotrophic bacterium *Klebsiella* sp. *Biol. Fertilizer of Soils* 1996;23(4):454-458.
10. Singh B, Singh YJ, Ladha KK, Bronson F, Balasubramanian V, Singh Y *et al.* Chlorophyll meter- and leaf color chart- based nitrogen management for paddy and wheat in northwestern India. *Agron. J*

2002;94:821-829.