www.ThePharmaJournal.com

The Pharma Innovation



ISSN (E): 2277-7695 ISSN (P): 2349-8242 NAAS Rating: 5.23 TPI 2022; SP-11(11): 323-326 © 2022 TPI www.thepharmajournal.com Received: 09-08-2022 Accepted: 15-09-2022

Affiliation all author is given at the end of references section

Determination of the optimum level of incorporation of exogenous fibrolytic enzymes, chicory root powder (*Cichorium intybus*) and their combination by *in vitro* experiment

JM Sanchaniya, HH Savsani, MD Odedra, JA Chavda, DD Garg, MR Chavda, DG Vaghamashi, NK Ribadiya, NJ Rankja, RB Makwana, VK Karangiya, HB Naliyapara, PH Agravat, YG Kansagara and SY Belim

Abstract

The present study was carried out with the objectives to determine the optimum level of incorporation of exogenous fibrolytic enzymes, chicory root powder and their combination by in vitro experiment. Sixteen healthy Gir calves of around one year of age were randomly selected from herd maintained at cattle breeding farm. Initially 2 Gir calves have been used to collect rumen liquor for in vitro study. Experimental calves were randomly divided on body weight basis into 4 groups having 4 calves in each group for in vivo study. The experimental feed was prepared using seasonal dry fodder as roughage and BIS type-1 compound cattle feed as concentrate in the ratio of 60:40. As a feed additive five different levels of exogenous fibrolytic enzymes (EFE) (0, 25000, 50000, 100000, 200000 IU per kg of concentrate) and five different levels of chicory root powder (CRP) (0, 0.5, 1, 2, 3% of concentrate) were used. The in vitro study was conducted to determine the In vitro Total Gas Production (IVTGP) with basal diet as substrate supplemented with various levels of fibrolytic enzyme, chicory root powder and combination to arrive at the optimum level of incorporation of these supplements for in vivo studies. The statistical analysis of data revealed significant effect of level of incorporation of different feed additives alongwith experimental feed on *in vitro* gas production. The result revealed significantly higher (p<0.01) IVTGP at 1% level of incorporation of CRP and 50,000 IU of EFE per kg of concentrate while in combination 1% CRP + 50000 IU of EFE showed statistically significant value of (p<0.05) IVTGP.

Keywords: Chicory root powder, exogenous fibrolytic enzymes, gir calves, in vitro dose optimization

1. Introduction

Gujarat is an important state in milk production and marketing in India based on co-operative dairy system. In India Gujarat is the 4th largest state in milk production based on co-operative dairy system. It contributed around 13.569 million tons of milk to the total milk pool of India Gir is an excellent dairy cattle breed, well-known among the milk producers for its milk producing ability, good fertility, heat tolerance, easy maintenance, resistance to diseases and longevity (NDDB, 2017)^[7]. Calves play an important role in development of the dairy sector of country, as the future of the dairy herd totally depends upon the successful raising of young calves (Anonymous, 2012)^[1].

One of the major constraints for the livestock population in India is qualitative and quantitative shortage of feed for livestock. The efficient conversion of energy-rich carbohydrates (cellulose, hemicellulose) of lignocellulosic biomass into accessible sugars is challenging technically, as these materials naturally evolved to resist degradation due to the complex fibrous structure of the materials that have constructed physical barriers to the accessibility of these carbohydrates for enzymatic breakdown (Thammiah *et al.*, 2017)^[13].

Cellulases and xylanases are the two major enzymes that are specified to break β 1-4 linkages joining sugar molecules of cellulose and xylans found in plant cell wall components (Beauchemin *et al.*, 2003)^[2].

The use of prebiotics is a promising approach for enhancing the role of endogenous beneficial microbiota in the gut (Patel and Goyal, 2012)^[8].

Corresponding Author: JM Sanchaniya Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry,

Kamdhenu University, Junagadh, Gujarat, India

Inulin (fructo-oligosaccharide) has positive effect in stimulation of the development of the gut microbiota strains, inhibition of the proliferation of pathogenic bacteria and helps in production of Short Chain Fatty Acids from the fermentation of dietary fiber also have significant impact on the immune system by acidification of the colonic environment, acidification of the colon will favors mucin production as well as binding to Short Chain Fatty Acid receptors on immune cells within the gut-associated lymphoid tissues (GALT) (Singh *et al.*, 2017) ^[10]. Keeping all these factors in the mind the *in vitro* experiment was carried out to optimize best level of incorporation of exogenous fibrolytic enzymes, chicory root powder and their combination for further *in vivo* study.

2. Materials and Methods

The present study was conducted on Sixteen Gir calves at Cattle Breeding Farm and Department of Animal Nutrition, College of Veterinary Science and A.H., Junagadh Agricultural University, Junagadh, Gujarat.

Rumen liquor was collected from two donor calves at 2 h post-feeding through a stomach tube against negative pressure created by a suction pump (Lane *et al.*, 1968)^[4]. The collected rumen liquor was strained through four layered muslin cloth and called as Strained Rumen Liquor (SRL). The SRL was brought to the laboratory in a pre-warmed ($39\pm1^{\circ}$ C) thermos flask. Carbon dioxide gas was passed through the SRL for one minute and was maintained at $39\pm1^{\circ}$ C temperature for further analysis.

The medium was prewarmed to 39 °C and bubbled with CO_2 till the blue color of the medium disappeared. After the medium becomes colorless strained rumen liquor (SRL) was added. Then 30 ml of incubation medium was injected to each calibrated glass syringe containing different levels of finely ground experimental feed along with fibrolytic enzymes (cellulase and xylanase), chicory root powder and combination as described by Menke *et al.* (1979) ^[6]. The experimental feed was prepared using seasonal dry fodder as roughage and BIS type-1 compound cattle feed as concentrate in the ratio of 60:40. As a feed additive five different levels of exogenous fibrolytic enzymes (EFE) (0, 25000, 50000, 100000, 200000 IU per kg of concentrate) and five different levels of chicory root powder (CRP) (0, 0.5, 1, 2, 3% of concentrate) was used.

The syringes were shaken gently, residual air or air bubble was removed and outlet was closed. The level of piston was recorded and the syringes were placed in an incubator $(39\pm1^{\circ}C)$ with periodic manual shaking (every 2 hour). Gas produced during fermentation was measured after 24 hrs. After incubation of 24 h, the content of each syringe was filtered through pre-weighed Gooch crucible. Composition if media for *in vitro* study is presented in Table 1.

The *in vitro* study was conducted to determine the *In vitro* Total Gas Production (IVTGP) with basal diet as substrate supplemented with various levels of fibrolytic enzyme, chicory root powder and combination to arrive at the optimum level of incorporation of these supplements for *in vivo* studies.

Statistical Analysis

The collected data was statistically analysed by two-way analysis of variance for *in vitro* study as per procedures suggested by Snedecor and Cochran (1994) ^[12]. The significance of mean differences was tested by Duncan's multiple range test (DMRT).

3. Results and Discussion

In vitro experiment were carried out using exogenous fibrolytic enzymes (cellulase and xylanase) and chicory root powder alone and in combination were used as feed supplement and added at different levels in the experimental feed which was prepared by using groundnut haulm as roughage and compound cattle feed type-1 as concentrate in the ratio of 60:40 to arrive optimum level of its incorporation by conducting *in vitro* experiment. Proximate composition for experimental feed is given in Table 2. The *in vitro* total gas production was studied to determine the optimum level of incorporation of exogenous fibrolytic enzymes (cellulase and xylanase) and chicory root powder as feed supplement.

In vitro total gas production (IVTGP) at 24 h incubation

The effect of individual feed additive as well as in combination at different levels of incorporation were assessed by *in vitro* gas production technique (Menke *et al.*, 1979)^[6]. The various levels used for exogenous fibrolytic enzymes and chicory root powder for *in vitro* study were (0, 25000, 50000, 100000, 200000 IU per kg of concentrate) and (0, 0.5, 1, 2, 3% of concentrate) respectively and the treatments were designated as EC_1 to EC_{25} (total 25 replications) for determination of optimum level of incorporation of exogenous fibrolytic enzyme and chicory root powder based on *in vitro* total gas production (TGP) profile.

In vitro gas production technique helps to determine nutrient utilization in a scientific way and its accuracy in describing digestibility in animals has been validated in numerous trials. This technique can be used to predict animal performance at much lower cost. Total gas production (ml) at 24 hours of incubation by *in vitro* gas production technique for Exogenous fibrolytic enzymes and Chicory root powder for different doses (total 25 combinations) using Gir calf rumen liquor to ascertain best level of exogenous fibrolytic enzymes, chicory root powder and best combination for further *in vivo* study have been presented in tabular form in Table 3.

The statistical analysis of data presented in table 4 revealed significant effect of level of incorporation of different feed additives alongwith experimental feed on *in vitro* gas production. The result revealed significantly higher (p <0.001) IVTGP at 1% level of incorporation of CRP and 50,000 IU of EFE per kg of concentrate while in combination 1% CRP + 50000 IU of EFE showed statistically significant value of (p < 0.005) IVTGP. However, data of *in vitro* trial indicates that beyond this level of incorporation resulted into reduction in IVTGP and no further beneficial effect were found.

The increase in total gas production appeared to be associated with higher DM and OM degradability in supplemented groups due to increase in bacterial number and activity. The rate of fermentation might have also increased in EFE and CRP supplemented groups and hence IVTGP was increased in present study.

Similar findings were also reported for EFE by many researchers, where higher IVTGP was observed due to enzyme supplementation. Elghndour *et al.* (2013)^[3], Sipai *et al.* (2013)^[11], Dey *et al.* (2014) and Lunagariya *et al.* (2017)^[5] who reported significant (p<0.05) increase in IVTGP by enzyme supplementation while Turner *et al.* (1999)^[14] and Salman *et al.* (2017)^[9] were also noticed similar trend of *in vitro* DM and OM degradability by incorporation of CRP/ inulin.

Table 1: Composition of media for in vitro study

Particulars	Quantity of different solutions and SRL (ml)		
Micro mineral solution	0.10		
Buffer solution	190.23		
Macro mineral solution	190.23		
Resazurin solution	0.95		
Reducing solution	38.05		
Distilled water	380.45		
Total media	800		
Rumen liquor	400		
Total mixture	1200		

 Table 2: Average proximate composition of feed/fodder for in vitro

 trial

Proximate	Ingredients (Feed/Fodder)		
Composition (% DM basis)	Groundnut haulms %	Compound cattle feed Type-1%	
DM	94	92	
OM	91.72	90.42	
СР	10.36	21.73	
EE	1.54	3.93	
CF	21.83	10.81	
NFE	57.99	53.95	
TA	8.28	9.58	

 Table 3: In vitro total gas production (ml/200mg experimental feed at 24 hrs.) under In vitro trial.

_	Level of Chicory root powder (CRP) (% of Concentrate)						
rate		0	0.5	1	2	3	
oncenti	0	EC 1 15	EC 2 18.5	EC 3 19**	EC 4 17	EC 5 14	
'el of EFE (IU/kg of Co	25,000	EC 6 16.5	EC7 20	EC8 17	EC 9 16.5	EC 10 17	
	50,000	EC 11 20.5**	EC 12 24	EC 13 23*	EC 14 18.5	EC 15 18.5	
	1,00,000	EC 16 19.5	EC 17 22.5	EC 18 20	EC 19 19	EC 20 17	
Lev	2,00,000	EC 21 18.5	EC 22 19.5	EC 23 17.5	EC 24 20	EC 25 15	

**= Significant at 1 per cent (p<0.01), *= Significant at 5 per cent (p<0.05)

Table 4: ANOVA of in vitro total gas production

Sources of variation	d.f.	MSS	F Value	P Value	Level of Significant
EFE	4	28.930	18.310	< 0.001	**
CRP	4	28.930	18.310	< 0.001	**
$EFE \times CRP$	16	3.343	2.116	0.045	*
Residual	25	1.580			
Total	49	6.621			
**- Significant at 10 (n < 0.01) *- Significant at 50 (n < 0.05)					

**= Significant at 1% (*p*<0.01), *= Significant at 5% (*p*<0.05)

4. Conclusions

Supplementation of exogenous fibrolytic enzymes (EFE) (50,000 IU/kg of concentrate), chicory root powder (CRP) (1% of concentrate) alone and their combination (EFE 50,000 IU/kg of concentrate + CRP 1% of concentrate) revealed Significantly higher (p < 0.001) *In vitro* total gas production (IVTGP).

5. Acknowledgement

Authors are thankful to Director of Research, Principle & Dean and Department of Animal Nutrition, College of Veterinary Science and A. H., Kamdhenu University, Junagadh, Gujarat for necessary facilities during research

work.

6. References

- Anonymous. 29th Survey Report on Estimates of Major Livestock Products for the year 2011-2012 Gujarat State, Directorate of Animal Husbandry Gujarat State, Krishibhavan, Sector-10/a, Gandhinagar; c2012. p. 114.
- 2. Beauchemin KA, Colombatto D, Morgavi DP, Yang WZ. Use of exogenous fibrolytic enzymes to improve feed utilization by ruminants. J Anim. Sci. 2003;81(2):37-47.
- 3. Elghandour MMY, Salem AZM, Gonzalez-Ronquillo M, Bórquez JL, Gado HM, Odongo NE, *et al.* Effects of exogenous enzymes on *in vitro* gas production kinetics and ruminal fermentation of four fibrous feeds. Anim. Feed Sci. Technol. 2013;179(1-4):46-53.
- 4. Lane GT, Noller CH, Clendrander VP, Cummings KR, Harzington RB. Apparatus for obtaining rumino-reticular samples and the effect of sampling location on pH and volatile fatty acids. J Dairy Sci. 1968;51(1):114-123.
- 5. Lunagariya PM, Gupta RS, Parnekar S. *In vitro* evaluation of total mixed ration supplemented with exogenous fibrolytic enzymes for crossbred cows. Vet. World. 2017;10(3):281-285.
- 6. Menke KH, Rabb L, Salewski A, Steingass H, Fritz D, Schneiser W. The estimation of the digestibility and metabolizable energy content of ruminant feedstuffs from the gas production when they are incubated with rumen liquor *in vitro*. J Agric. Sci. 1979;93(1):217-222.
- NDDB. NDDB Report Entitled Approaches and experiences of NDDB in development of Gir a promising indigenous milch breed report downloaded on 20th December 2017-2019 from https://www.nddb.coop/sites/default/files/Booklet-Development-of-Gir-Eng% 20-low.pdf Pp. 36.
- 8. Patel S, Goyal A. The current trends and future perspectives of prebiotics research: a review. Biotech. 2012;2(2):115-125.
- Salman M, Cetinkaya N, Selcuk Z, Genc, Acici M. Effects of various inulin levels on *in vitro* digestibility of corn silage, perennial ryegrass (*Lolium perenne* L.) and common vetch (*Vicia sativa* L.)/oat (*Avena sativa* L.) hay. S. Afr. J Anim. Sci. 2017;47(5):723-729.
- Singh AK, Kerketta S, Yogi RK, Kumar A, Ojha L. Prebiotics-The new feed supplement for dairy calf. Int. J Livest. Res. 2017;7(8):1-17.
- 11. Sipai SH, Dutta KS, Savsani HH, Murthy KS, Vataliya PH, Chavda JA, *et al.* Inclusion of different exogenous fibrolytic enzymes to dry jowar fodder and their effect on *in vitro* total gas production. Vet. World. 2013;6(11):839-843.
- 12. Snedcor G, Cochran W. Statistical Methods. 8th Edition. Oxford and IBH. New Delhi; c1994.
- 13. Thammiah V, Samanta AK, Senani S, Manpal S. Scope of exogenous enzymes in enhancing ruminant productivity. J Dairy. Vet. Anim. Res. 2017;5(2):67-72.
- 14. Turner KE, Belesky DP, Fedders JM. Chicory effects on lamb weight gain and rate of *in vitro* organic matter and fibre disappearance. Agron. J. 1999;91(3):445-450.

Authors Details

JM Sanchaniya

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

HH Savsani

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

MD Odedra

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

JA Chavda

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

DD Garg

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

MR Chavda

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh

DG Vaghamashi

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

K Ribadiya

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

NJ Rankja

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

RB Makwana

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

VK Karangiya

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

HB Naliyapara

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

PH Agravat

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India

YG Kansagara

Department of Animal Nutrition, College of Veterinary

SY Belim

Department of Animal Nutrition, College of Veterinary Science and Animal Husbandry, Kamdhenu University, Junagadh, Gujarat, India