



ISSN (E): 2277-7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2022; SP-11(11): 2021-2024
© 2022 TPI
www.thepharmajournal.com
Received: 28-09-2022
Accepted: 30-10-2022

Insha Afzal
Division of L.P.M, FVSc & AH,
SKUAST K, Shuhama Alusteng,
Ganderbal, Jammu and
Kashmir, India

Azmat Alam Khan
Division of L.P.M, FVSc & AH,
SKUAST K, Shuhama Alusteng,
Ganderbal, Jammu and
Kashmir, India

Sofi Umer Bashir
Department of Wildlife Science,
Madras Veterinary College,
Tamil Nadu Veterinary and
Animal Sciences University,
Chennai, Tamil Nadu, India

Ashaq Manzoor
Division of L.P.M, FVSc & AH,
SKUAST K, Shuhama Alusteng,
Ganderbal, Jammu and
Kashmir, India

Impact of bovine colostrum used as feed additive in broiler chicken diets on body weight, mortality pattern and economics of production

Insha Afzal, Azmat Alam Khan, Sofi Umer Bashir and Ashaq Manzoor

Abstract

A biological trial was conducted on 192, three days old commercial broiler chicks, distributed randomly into four treatment groups of 4 replicates of 12 birds each, reared for a period of 35 days. The first treatment group (T₁) was taken as control group and offered basal diet without any additive. The other three treatment groups (T₂, T₃ and T₄) were fed diets supplemented with 1, 1.5 and 2 % bovine colostrum/kg feed, respectively. The treatment groups offered diets supplemented with bovine colostrum showed significantly higher body weight with the highest body weight recorded in T₄. The final body weights recorded at the end of the trial period were 1503.04±25.16, 1577.99±35.99, 1675.86±77.21 and 1897.02±37.51 in groups T₁, T₂, T₃ and T₄ respectively. There was a significant reduction in mortality percentage across the treatment groups. An overall mortality of 11.43% was recorded during the trial. The highest overall mortality being recorded in T₁ (19.23%) and lowest in T₄ (2.5%). T₂ and T₃ showed a mortality percentage of 16.92 and 3.77 respectively. There was a positive impact on the economics of production, thus increasing the profit margins. Economic analysis revealed that the cost of feed/kg body weight gain was Rs 78.20, 73.50, 69.58 and 66.60 in T₁, T₂, T₃ and T₄ respectively. T₂, T₃ and T₄ had a relative advantage of Rs 4.70, 8.62 and 11.60 respectively as compared to control T₁. Thus, it was concluded that bovine colostrum could be incorporated in broiler diets @ 2% for better body weight at market age, reduced mortality percentage and improved the overall economics of production.

Keywords: Broilers, bovine colostrum, body weight, mortality, economics

1. Introduction

The Poultry industry has been recognized as the fastest growing segment of the Indian agricultural sector. Most of this growth could be attributed to commercial broiler and layer enterprises that have come up since early eighties. The broiler industry in particular is growing very fast. The current strength of broilers in India is estimated to be 2300 million with the broiler meat production of 2.6 million tons thus making India the 5th largest poultry meat producer in world [24].

Poultry production, particularly broiler production is the quickest way to increase the availability of high quality protein for human consumption. Since the feed cost alone contributes to about 70-75% of the total cost of production, economically poultry production is, therefore, possible only when the feed cost is reduced and efficiency of feed utilization is increased [17]. Several studies have shown that use of feed additives considerably reduces feed cost and economises production in broiler chicken either by improving body weight, FCR or both and decreasing mortality percentage. Additionally, the efforts are on to identify and use various feed additives/ nutraceuticals/ bio-active compounds to optimize nutrient utilization, by favorably altering the intestinal microflora and intestinal micro architecture and by enhancing the immunity of the birds so as to prevent the losses on account of morbidity and mortality.

Colostrum is the “early” milk produced by female mammals and in case of bovines it is the first four days of milk post parturition [7]. In addition to proteins, carbohydrates, fats, vitamins and minerals it contains various growth factors, cytokines and nucleosides. It is rich in oligosaccharides, natural antimicrobials, immune regulation factors [9] and anti-oxidative factors. Bovine colostrum contains immuno-globulins such as IgG, IgM, IgA, IgD, and IgE. IgG and IgM function in systemic infections while IgA functions within internal body surfaces such as the intestine [27]. When taken orally, the immuno-globulins in colostrum may not be absorbed into the blood stream. However, within the gastro-intestinal tract of animals of all ages they are very effective against pathogenic organisms.

Corresponding Author:
Insha Afzal
Division of L.P.M, FVSc & AH,
SKUAST K, Shuhama Alusteng,
Ganderbal, Jammu and
Kashmir, India

Colostrum immunoglobulins are able to bind and agglutinate invading organisms such as bacteria, viruses, fungi and parasites that enter the intestinal tract, facilitating their removal before they cause infection and diseases [25]. The immunoglobulins also prevent pathogens from binding to intestinal surfaces [26] thereby inhibiting an important step in the infection process. Published reports have shown that colostrum and its components are effective against a wide range of common pathogens which include *Rotavirus*, *Cryptosporidium* spp., *Staphylococcus aureus*, *Candida* spp., *Clostridium* spp., *Feline immunodeficiency virus*, *Shigella* spp., *Streptococcus* spp. and *E. coli*.

Keeping in view the versatility of colostrum along with its immense food, nutritional and economic value, the present study was envisaged to study the effect of incorporation of bovine colostrum in broiler rations on body weight, mortality percentage and economics of production.

2. Materials and methods

Bovine Colostrum from Six Crossbred Jersey Cows secreted immediately after calving was procured and stored in sterile containers in packs of 250 ml or less and frozen immediately till use. A biological trial duly approved by the Institutional Animal Ethics Committee (IAEC) was conducted on one hundred and ninety two, three days old commercial broiler chicks (Rathi cob) procured from a reputed hatchery and reared for a period of 36 days. The birds were reared on deep litter throughout the experimental period. The chicks were distributed randomly into four treatment groups (T₁, T₂, T₃ and T₄) and each treatment comprised of 4 replicates of 12 birds each. Group T₁ was offered basal diet comprising of commercially available broiler pre-starter and starter rations while T₂, T₃ and T₄ were offered basal diets supplemented with 1, 1.5 and 2% of colostrum/kg feed. All chicks were kept under the same managerial, hygienic and environmental conditions and had *ad libitum* access to feed and water throughout and were maintained on a constant 24 hours light schedule. All chicks were vaccinated against Ranikhet disease on the 5th day with F₁ strain vaccine and B₂K vaccine against Infectious Bursal Disease (IBD) on the 16th day.

2.1 Live body weight

The body weight of the experimental birds was recorded on the individual basis at weekly intervals.

2.2 Mortality

The chicks were checked twice daily for mortality, if any and the same was recorded.

2.3 Economics of production

The economics of production at the end of the experimental period were calculated using market prices of feed, colostrum, feed consumed and body weight gain.

2.4 Statistical Analysis

The analysis of the data was carried out by using Analysis of Variance (ANOVA). The data generated was classified treatment wise and averages were drawn. After performing statistical analysis the tests were referred by p-values. Any p-value less than 0.05 ($p < 0.05$) was taken as statistically significant. The comprehensive statistical package for social sciences (SPSS ver.15.00) Chicago U.S.A. for Windows was

used for analysis.

3. Results

3.1 Body Weight

The results of weekly live body weight of broiler chicken under different dietary treatments have been summarised in Table 1. The body weight after 1 week of feeding bovine colostrum (BC) incorporated diets showed an increasing trend from T₁ to T₄ but there was no significant difference between the groups. From 2nd week onwards, there was a gradual increase in body weight of colostrum fed groups compared to control, with significantly higher live body weight in the group fed 2% BC in the diet. At the end of 3rd week also, the group supplemented with 2% BC showed significantly higher body weight as compared to control group and also in comparison to other BC treated groups. Similarly at the end of 4th week, the group fed 2% BC supplemented ration showed significant increase in body weight as compared to the control group but no significant difference in body weight was noticed in comparison with other BC supplemented groups. The final body weight of the birds at the end of five weeks of feeding trial showed an increasing trend from T₁ to T₄ groups. Final body weight was significantly higher in the groups fed BC supplemented feed as compared to control. 2% BC supplemented group had significantly higher body weight than 1% and 1.5% BC supplemented groups as well.

3.2 Mortality

The weekly and overall mortality of the broiler birds subjected to different concentrations of BC in the feed is depicted in Table 2. An overall mortality of 11.43% was recorded during the trial. An increasing trend was seen in mortality in different weeks with a mortality of 1.90% in 1st week, 2.43% in 2nd week, 2.49% in 3rd week, 2.04% in 4th week and 3.13% in 5th week was recorded. The overall mortality across the groups however showed a decreasing trend with the highest overall mortality of 19.23% recorded in group T₁ and lowest of 2.5% in group T₄. Treatment groups T₂ and T₃ showed a mortality percentage of 16.92 and 3.77 respectively. Between the groups higher mortality was recorded in 5th week in group T₂ (5.26%) followed by T₁ (4.55%) and least in T₃ (1.92%). T₄ had 0% mortality during this period. The study revealed that BC supplementation resulted in lower mortality in birds and it was considerably lower in group T₄ supplemented with the highest percentage of BC(2%).

3.3 Economics

The economics of production of broiler chicken fed BC supplemented diet in different concentrations worked out on the basis of feed cost per kg live weight gain is depicted in Table 3. The feed cost/kg body weight gain was Rs 78.20, 73.50, 69.58 and 66.60 in groups T₁, T₂, T₃ and T₄ respectively. There was a relative advantage of Rs 4.70, 8.62, 11.60 per kilogram of live weight in groups T₂, T₃ and T₄ respectively as compared to the control group T₁. The economics worked out at the end of the trial based on feed cost/kg live weight indicated higher feed cost in the group not supplemented with BC (T₁) and least in the group supplemented with 2% BC (T₄). Feed cost/kg live weight thus showed a decreasing trend across the groups.

Table 1: Effect of incorporation of Bovine colostrum in broiler chicken diet on Body weight and Body weight gain

Weeks of supplementation	Treatment groups			
	T ₁ (0 % colostrum)	T ₂ (1 % colostrum)	T ₃ (1.5 % colostrum)	T ₄ (2 % colostrum)
Body weight (g)				
0 day	96.29±1.04	94.03±1.12	96.21±1.00	96.92±0.82
1	274.69±2.90	274.34±1.98	276.16±4.53	280.74±5.13
2	534.37±10.22 ^a	530.89±9.96 ^a	550.96±5.11 ^{ab}	572.65±13.29 ^b
3	663.28±5.52 ^a	704.34±8.54 ^b	721.68±11.69 ^b	766.73±21.26 ^c
4	992.50±22.46 ^a	1085.47±57.16 ^{ab}	1140.56±63.68 ^{ab}	1201.92±100.52 ^b
5	1503.04±25.16 ^a	1577.99±35.99 ^{ab}	1675.86±77.21 ^b	1897.02±37.51 ^c

Means within the same row with different superscripts are significantly different ($p \leq 0.05$)

Table 2: Effect of incorporation of Bovine colostrum in broiler chicken diet on mortality (%)

Age in weeks	Treatment groups				Overall
	T ₁ (0% colostrum)	T ₂ (1% colostrum)	T ₃ (1.5% colostrum)	T ₄ (2% colostrum)	
0-1	3.85	3.08	0.00	0.00	1.90
1-2	4.00	1.59	1.89	2.55	2.43
2-3	4.17	4.84	0.00	0.00	2.49
3-4	4.35	3.39	0.00	0.00	2.04
4-5	4.55	5.26	1.92	0.00	3.13
0-5	19.23	16.92	3.77	2.5	11.43

Table 3: Effect of incorporation of Bovine Colostrum in broiler chicken diet on Economics of production

	Treatment groups			
	T ₁ (0% colostrum)	T ₂ (1% colostrum)	T ₃ (1.5% colostrum)	T ₄ (2% colostrum)
FCR	2.30	2.10	1.96	1.85
Cost of Basal Feed (Rs)/Kg	34	34	34	34
Cost of Colostrum (Rs)/Kg of feed	0	1.00	1.50	2.00
Total Cost of Feed (Rs/Kg)	34	35.00	35.50	36.00
Feed Cost/Kg live wt gain (Rs)	78.20	73.50	69.58	66.60
Difference in Feed cost/Kg live wt gain (Rs) as compared to control	-	4.70	8.62	11.60

4. Discussion

In the present study, significant ($p \leq 0.05$) difference in body weight was observed in the group supplemented with 2% BC (T₄). A similar study was conducted which resulted in higher body weight at day 7 and 13 in immunmilk fed birds as compared to the control^[18]. Also a patent, that stands filed at US patent office with regards to use of a feed additive containing bovine colostrum claims that incorporation of this feed additive enhances growth in broiler chicken^[3]. Some of the growth promoting proteins that have been identified in colostrum include insulin-like growth factors (IGF) 1 and 2^[6, 22], transforming growth factor (TGF-B)^[15], platelet derived growth factor (PDGF) like molecule termed as colostrum growth factor^[21], and angionin^[4]. It is therefore possible that the colostrum improves growth and integrity of the intestinal tract thereby allowing better nutrient absorption and utilization resulting in higher body weight and body weight gain^[18]. However, there was no effect of feeding spray-dried colostrum on body weight of chick^[10].

The results with respect to mortality revealed that BC supplementation resulted in lower mortality in birds and it was considerably lower in group T₄ supplemented with highest percentage of BC(2%). Our results are in agreement with those of other workers who found similar results in other animals. An increased survival rate in weaned piglets was reported on BC supplementation^[23]. A better steady-state of red kids after supplemental intake of colostrum has also been reported by some workers^[1, 13]. This may be directly associated with the presence of various nutrients in colostrum. Besides providing energy and protein, the supplementation of colostrum probably helped bringing additional levels of vitamins (A and E) and minerals (zinc, selenium and iodine)

which gave animals a way to develop and maintain a stronger immunity^[12, 20]. Colostrum is also rich in lactoferrin, which plays an important role in defending the body against pathogens^[2, 5, 20]. In addition, colostrum provides at low concentration polypeptidic compounds such as cytokines (IL-1, IL-6, IFN-7 and IL-1) belonging to the family of specific and non-specific antimicrobial factors^[8].

The results of economics worked out at the end of the trial based on feed cost/kg live weight indicated higher feed cost in the group not supplemented with BC (T₁) and least in the group supplemented with 2% BC (T₄). This could be a direct consequence of increased weight gain and improved FCR noted in the groups subjected to BC supplementation. Thus, addition of feed additives to the broiler diets economises the production cost as investigated by various workers. The effect of six commonly available commercial feed additives on the performance of the broilers was studied and the ration having no feed additive (control) was found to be the most uneconomical^[16]. Similarly, the addition of commercial and natural (skimmed milk and whey) probiotics to broiler diets improved the average values of calculated economical efficiency^[14]. Addition of phytochemicals (Dandelion leaves and fenugreek seeds) to the broiler diet also resulted in reduction in feed cost/kg live weight as compared to the control^[19]. The diet supplemented with whey protein was cheaper than the control and the highest profitability was recorded with the broiler chicks which received the highest percentage of whey protein among the various groups^[11].

5. Conclusion

It was concluded that adding 2% bovine colostrum to broiler chicken diets had a positive impact on body weight of broiler

chicken and reduced mortality percentage thereby resulting in improved overall economics of production.

6. Acknowledgement

Division of Livestock Production and Management, FVSc and AH, SKUAST-K, Shuhama, Kashmir.

7. References

- Abdou H, Marichatou H, Beckers JF, Dufrasme I, Issa M, Homick JL. Effect of bovine colostrum intake on growth, reproductive parameters and survival in red kids. *Journal of Animal Physiology and Animal Nutrition* 2014;98:845-852.
- Ajello M, Greco R, Giansanti F, Massucci MT, Antonini G, Valenti P. Anti-invasive activity of bovine lactoferrin towards group A streptococci. *Biochemistry and Cell Biology*. 2002;80:119-124.
- Anonymous. Organic/natural poultry feed additive Patent No: US 20090226558 A1 filed by C and R BIOTEK, LLC, GEORGIA 2009; <http://www.google.com/patents/US20090226558>.
- Chang SI, Jeong GB, Park SH, Ahn BC, Choi JD, Chae Q, *et al.* Detection, quantitation, and localization of bovine angiogenin by immunological assays. *Biochemical and Biophysical Research Communications*. 1997;232:323-327.
- Di Mario F, Aragona G, Bò ND, Ingegnoli A, Cavestro GM, Moussa AM, *et al.* Use of lacto-ferrin for *Helicobacter pylori* eradication. Preliminary results. *Journal of Clinical Gastroenterology*. 2003;36:396-398.
- Francis GL, Upton FM, Ballard FJ, McNeil KA, Wallace JC. Insulin-like growth factors 1 and 2 in bovine colostrum. Sequences and biological activities compared with those of a potent truncated form. *Biochemical Journal*. 1988;251:95-103.
- Gopal PK, Gill HS. Oligosaccharides and glycoconjugates in bovine milk and colostrum. *British Journal of Nutrition*. 2000;84(S1):69-74.
- Hagiwara K, Kataoka S, Yamanaka H, Kirisawa R, Iwai H. Detection of cytokines in bovine colostrum. *Veterinary Immunology Immunopathology*. 2000;76:183-190.
- Kelly GS. Bovine colostrums: A review of clinical uses. *Alternate Medicine Reviews*. 2003;8:378-394.
- King MR, Ravindran V, Morel PCH, Thomas DV, Birtles MJ, Pluske JR. Effects of spray-dried colostrum and plasmas on the performance and gut morphology of broiler chickens. *Australian Journal of Agricultural Research*. 2005;56:811-817.
- Malik HEE, Elamin KM, Abdalla SA, Dousa BM. Influence of Supplemented Whey On Growth Performance And Internal Organs Percentages Of Broiler Chicken. *Online Journal of Animal and Feed Research*. 2015;5(3):68-73.
- Morales-delaNuez A, Moreno-Indias I, Sánchez-Macías D, Capote J, Juste MC, Castro N, *et al.* Sodium dodecyl sulfate reduces bacterial contamination in goatcolostrum without negative effects on immune passive transfer in goat kids. *Journal of Dairy Science*. 2011;94:410-415.
- Moretti DB, Kindlein L, Pauletti P, Machado-Neto R. IgG absorption by santa Ines lambs fed Holstein bovine colostrum or Santa Inesovine colostrum. *Animal*. 2010;6:933-937.
- Omara II. Nutritive value of skimmed milk and whey, added as natural probiotics in broiler diets. *Egyptian Journal of Animal Production*. 2012;49:207-217.
- Pakkanen R. Determination of transforming growth factor-beta 2 (TGF-beta 2) in bovine colostrum samples. *Journal of Immunoassay*. 1998;19:23-37.
- Pervez, Rafiullahand Sajid A. Effect of Feed Additives on the performance of broilers. *ARNP Journal of Agricultural and Biological Science*. 2011;6:9.
- Qureshi AA. Selecting the best feeding and watering equipment. *Misset World Poultry*. 1991;7:17-19.
- Qureshi MA, Ali R, Cheema MA, Ahmed Z, Roth H. Immunmilk Feeding Increases Growth and Immune Responses in Broiler Chicks. *International Journal of Poultry Science*. 2004;3(5):305-312.
- Qureshi S. Evaluation of locally available herbs Dandelion leaves (*Taraxacum officinale*) and Fenugreek seeds (*Trigonella foenum-graceum*) as feed additives in Broiler Chicken ration 2014; M.V.Sc Thesis submitted to SKUAST-K, Shalimar.
- Rawal P, Gupta V, Thapa BR. Role of Colostrum in Gastrointestinal Infections. *Indian Journal of Pediatrics*. 2008;75(9):917-921.
- Shing Y, Klagsbrun M. Purification and characterization of a bovine colostrum-derived growth factor. *Molecular Endocrinology*. 1987;1:335-338.
- Vega JR, Gibson CA, Skaar TC, Hadsell DL, Baumrucker CR. Insulin-like growth factor (IGF)-I and -II and IGF binding proteins in serum and mammary secretions during the dry period and early lactation in dairy cows. *Journal of Animal Science*. 1991;69:2538-2547.
- Viehmann M, Unterweger C, Ganter M, Metzler-Zebeli BU, Ritzmann M, Hennig-Pauka I. Effects of bovine colostrum on performance, survival, and immunoglobulin status of suckling piglets during the first days of life. *Czech Journal of Animal Science*. 2015;60(8):351-358.
- Anonymous 2014. Poultry sector in India. Poultry Fest, 3rd edition. <http://www.poultryfest.in/poultryfest.php>.2014.
- Oyenyi OO, Hunte AG. Colostral constituents including immunoglobulins in the first three milkings postpartum. *Journal of Dairy Science*. 1978;61:44-48.
- Bitzan MM, Gold BD, PhilPott DJ. Inhibition of *Helicobacter pylori* and *Helicobacter mustelae* binding to lipid receptors by bovine colostrum. *Journal of Infectious Disease*. 1998;177:955-961.
- Muller LD, Ellinger DK. Colostral immunoglobulin concentrations among breeds of dairy cattle. *Journal of Dairy Science*. 1981;64:1727-1730.