



ISSN (E): 2277- 7695

ISSN (P): 2349-8242

NAAS Rating: 5.23

TPI 2022; 11(2): 09-12

© 2022 TPI

www.thepharmajournal.com

Received: 07-11-2021

Accepted: 16-12-2021

Poonam Kumari

Department of Plant Pathology,
Rajasthan College of Agriculture,
MPUAT, Udaipur, Rajasthan,
India

Amit Trivedi

Department of Extension
Education, Rajasthan College of
Agriculture, MPUAT, Udaipur,
Rajasthan, India

Ranjana Meena

Department of Plant Pathology,
Sri Karan Narendra Agriculture
University, Jobner, Jaipur,
Rajasthan, India

Pinki Devi Yadav

Department of Plant Pathology,
Sri Karan Narendra Agriculture
University, Jobner, Jaipur,
Rajasthan, India

Monika Meena

Department of Plant Pathology,
Sri Karan Narendra Agriculture
University, Jobner, Jaipur,
Rajasthan, India

Sunita Choudhary

Department of Plant Pathology,
Sri Karan Narendra Agriculture
University, Jobner, Jaipur,
Rajasthan, India

Corresponding Author:

Poonam Kumari

Department of Plant Pathology,
Rajasthan College of Agriculture,
MPUAT, Udaipur, Rajasthan,
India

Evaluation of different fungicides, combi products and botanicals efficacy against *Alternaria alternata* on Asalio in control condition

Poonam Kumari, Amit Trivedi, Ranjana Meena, Pinki Devi Yadav, Monika Meena and Sunita Choudhary

Abstract

Alternaria, the fungal pathogen has wide host range generally attacks the aerial parts of plants causing leaf spots. Among the different diseases affecting Asalio, leaf blight caused by *Alternaria alternata* is an important one. Very little information is available on the management aspects of this disease. Therefore, we evaluate two combi products, two individual fungicides and two botanicals was evaluated on potato-dextrose-agar media for sensitivity against *A. alternata*. Among these treatments combi products *i.e.*, Metiram 70% + Pyraclostrobin 20% WG (0.35% concentration) was found most effective in inhibiting mycelial growth of *A. alternata* *In vitro*.

Keywords: Per cent disease index, *Lepidium sativum*, *Alternaria alternata*, *In vitro*, botanicals, Combi products and fungicides

Introduction

Asalio (*Lepidium sativum* Linn. Family: Brassicaceae) is a medicinal plant. *Lepidium* name derives from Greek word 'lepidion' means small scale probably it refers to the form of fruits and *sativum* is derived from 'serere' meaning to cultivate, to plant or to sow. It is known as "Common cress", "Land cress", "Haliv", "Garden cress" or "Chandrasur" in some regions of India (Gokavi *et al.*, 2004) [3].

The crop is mainly affected by diseases like *Alternaria* leaf spot & white rust. The Rabi season Asalio crop in India is commonly affected by *Alternaria* leaf spot. The disease causes extensive damage to the quality and quantity of the foliage and grains. If persistent cloudy and cool weather prevails the blight appears after flowering stage of the crop. *Alternaria* genus is the largest genus. It is distributed worldwide and has been reported on 115 plant genera from 43 plant families that cause blight and leaf spot disease (Neergaard, 1945) [8]. Its species have wide host range fungi and it is causing crop loss in economically important crop plants by causing leaf spot diseases. *Alternaria* pathogen have a large host range, in terms of total number of host plants it has 10th rank in all the pathogens (Farr *et al.*, 1989) [2]. *Alternaria* genus causes different diseases, in which leaf spot disease is most severe to the crop and losses to the tune of 32-57 per cent in yield on a wide range of crops (Conn and Tewari, 1990) [1]. In some parts of the world It is reported to have has 4 per cent fungal diversity and have been reported to cause losses up to 80 per cent (Rotem, 1994) [9].

There is a need to identify efficient management strategies of this disease. *In vitro* evaluation of fungicides helps to identify effective fungicide for application in fields. In the present investigation an attempt was made to identify effective fungicides to manage *A. alternata*.

Materials and Methods

In vitro evaluation of different fungicides and botanicals was carried out using poisoned food technique. Test Fungicides and botanicals as per the treatments were evaluated on potato dextrose agar (PDA) medium. The flasks containing melted and sterilized PDA was amended with the desired concentration of each test fungicide and botanical. 20 ml of the amended PDA was poured in 90 mm sterilized Petri-plates and allowed to solidify. From the periphery of 7-days-old actively growing and sporulating culture of *A. alternata*, 5 mm disc was cut and placed in the centre of each experimental Petri-plate under totally aseptic conditions. The experiment was conducted in completely randomized design (CRD) with five replications in each treatment. These were then incubated at 27±1 °C in BOD. After 7 days when the control plates were full of fungal growth, the radial diameter was measured diagonally in each of the control and treatment plates.

Table 1: Treatment details

Number	Name of treatments	Concentrations
T1	Azoxystrobin 18.2% w/w + Difenoconazole 11.4% w/w SC	0.07%
T2	Azoxystrobin 23 SC	0.1%
T3	Difenoconazole 25 EC	0.06%
T4	Metiram 70% + Pyraclostrobin 20% WG	0.35%
T5	Azadirachtin	0.5%
T6	Neem oil	0.5%
T7	Control	

Design-CRD, Replication-5, Treatment: 7

Per cent inhibition of mycelial growth was calculated by using formula given by Bliss (1934) as:

$$I = \frac{C-T}{C} \times 100$$

Where

I = Per cent inhibition

C = Colony diameter in control

T = Colony diameter in treatment

Results and Discussion

In vitro bioassay of different fungicides and botanicals in amended potato dextrose agar media against *Alternaria alternata*

Efficacy of different fungicides and botanicals *viz.*, Azoxystrobin 18.2% w/w + Difenoconazole 11.4% w/w SC (0.07% concentration), Azoxystrobin 23 SC – (0.1% concentration), Difenoconazole 25 EC – (0.06% concentration), Metiram 70% + Pyraclostrobin 20% WG – (0.35% concentration), Azadirachtin – (0.5% concentration), and Neem oil – (0.5% concentration), (as discussed in Materials and Methods) were tested *in vitro* against mycelial growth of *A. alternata* by Poisoned Food Technique. The Results envisage that the combi products Metiram 70% +

Pyraclostrobin 20% WG –(0.35% concentration) with the colony diameter 18.64 mm followed by Azoxystrobin 18.2% w/w + Difenoconazole 11.4% w/w SC – (0.07% concentration), with colony diameter 28.04 mm at given concentrations were significantly superior in inhibiting the mycelial growth of the test fungus over the individual chemicals and control. Among the individual chemicals Difenoconazole 25 EC (0.06%concentration) with colony diameter 36.48 mm was significantly superior in inhibiting the growth of test fungus followed by Azoxystrobin 23 SC (0.1% concentration) with colony diameter 38.51 mm. All the test chemicals as combi products or individually were found to inhibit the growth of *A. alternata* significantly over botanicals and control. As for botanicals Neem oil –(0.5% concentration) was significantly more effective within test botanicals with colony diameter 45.90 mm then Azadirachtin – (0.5% concentration) with colony diameter 65.84 mm in inhibiting mycelial growth of *A. alternata*.

The Results reveal that the combi products Metiram 70% + Pyraclostrobin 20% WG – (0.35% concentration) with the per cent growth inhibition 79.28 followed by Azoxystrobin 18.2% w/w + Difenoconazole 11.4% w/w SC – (0.07% concentration), with the per cent growth inhibition 68.84 at given concentrations were significantly superior in inhibiting the growth of the test fungus over the individual chemicals and control. Among the individual chemicals Difenoconazole 25 EC (0.06% concentration) with the per cent growth inhibition 59.24 was significantly superior in inhibiting the growth of test fungus followed by Azoxystrobin 23 SC (0.1% concentration) with the per cent growth inhibition 57.21 mm. Among botanicals tested Neem oil – (0.5% concentration) was significantly more effective within test botanicals with the per cent growth inhibition 49.00 then Azadirachtin – (0.5% concentration) with the per cent growth inhibition 26.84 in inhibiting the growth of *A. alternata* (Table-1, Plate-5 and Fig-1).

Table 2: *In vitro* efficacy of different fungicides and botanicals against *Alternaria alternata* after 7 days inoculation at 27±1 °C

S. No.	Treatments	Colony diameter(mm)*	Per cent growth inhibition**
1	Azoxystrobin 18.2% w/w +Difenoconazole 11.4% w/w SC (0.07)	28.04	68.84 (56.08)
2	Azoxystrobin 23 SC (0.1)	38.51	57.21 (49.13)
3	Difenoconazole 25 EC (0.06)	36.48	59.24 (50.36)
4	Metiram 70% + Pyraclostrobin 20% WG (0.35)	18.64	79.28 (62.90)
5	Azadirachtin (0.5)	65.84	26.84 (31.18)
6	Neem oil (0.5)	45.90	49.00 (44.41)
7	Control	90.00	0.00
	S.Em±	1.466	1.636
	CD at 5%	4.268	4.763
	CV %	7.090	7.522

* Mean of five replications

Figures in parentheses are arcsine√per cent angular transformed values.




Fig 1: *In vitro* evaluation of different fungicides and botanicals against *A. alternata*



T1 Azoxystrobin 18.2% w/w + Difenconazole 11.4% w/w SC (0.07)
 T2 Azoxystrobin 23 SC (0.1)
 T3 Difenconazole 25 EC (0.06)
 T4 Metiram 70% + Pyraclostrobin 20% WG (0.35)
 T5 Azadirachtin (0.5)
 T6 Neem oil (0.5)
 T7 Control

Fig 2: *In vitro* evaluation of different fungicides and botanicals against *Alternaria alternata* after 7 days at 27 + 1 °C

As narrated earlier there have been no reports in the review of literature for management of *A. alternata* causing Alternaria leaf spot of Asalio. Although, management of *A. alternata* causing leaf spots in other crops is listed here. Survilienė and Dambrauskienė (2006) [10] reported efficacy of Azoxystrobin 250 SC against various *Alternaria* species viz., *Alternaria alternata*, *Alternaria brassicae*, *Alternaria dauci* on potato dextrose agar under *In vitro* conditions and reported 87% to

96% inhibition of colonies of different *Alternaria* species by Azoxystrobin at 7 days after inoculation. Mane (2008) reported that Mancozeb (0.2%) and Propiconazole (0.05%) completely inhibited the growth of *A. alternata* causing leaf blight of chilli. Kamanna *et al.* (2010) [4] observed that Chlorothalonil (0.2%) was effective for the control of chrysanthemum leaf blight under field conditions caused by *A. alternata*. The present study are liken with the farther said

reported reviews.

In view of the increasing disease incidence, in Asalio growing areas, attempts were made to evaluate two combi products and two fungicides and two botanicals formulations against *A. alternata*, which were first tested *In vitro*, combi products Metiram 70% + Pyraclostrobin 20% WG -0.35% and Azoxystrobin 18.2% w/w + Difenconazole 11.4% w/w SC -0.07%, were most effective in inhibiting growth of *A. alternata* *In vitro* with per cent inhibition of growth as 79.28% and 68.84%, respectively. Azadirachtin has been reported to be safe for the environment (Khalid and Shad, 2002; Morgan, 2009) but no information is available on efficacy of botanicals in suppressing *A. alternata* causing Alternaria leaf spot disease of Asalio. Among botanicals Neem oil effectively inhibited the growth of *A. alternata* in present study with per cent growth inhibition as 49.00%. When treated with fungicides, botanicals and combi products showed variations in suppressing *A. alternata*. The observations suggest that sensitivity to combi products is greatly prevalent in natural population of *A. alternata*.

References

1. Conn KL, Tewari JP. Survey of Alternaria black spot and Sclerotinia stem rot in central Alberta in 1989. Can. J Plant Dis. Survey. 1990;70:66-67.
2. Farr DF, Bills GF, Chamuris GP, Rossman AY. Fungi on Plants and Plant Products in the United States. The American Phytopathological Society press, St. Paul, (Minnesota) USA. 1989.
3. Gokavi SS, Malleshi NG, Guo M. Chemical composition of garden cress (*Lepidium sativum*) seeds and its fractions and use of bran as a functional ingredient. Journal of Plant Food and Human Nutrition. 2004;59:105-111.
4. Kamanna BC, Shankarappa TH, Arunkumar GS. Evaluation of fungicides for the management of chrysanthemum leaf blight caused by *Alternaria alternata* (Fr.) Keissler. Plant Archives. 2010;10:595-597.
5. Khalid S, Shad RA. Potential advantage of recent allelochemical discoveries and agro-ecosystems. Prog. Farm. 2002;11:30-35.
6. Mane VA. Studies on Alternaria blight and Fusarium wilt diseases of chilli (*Capsicum annuum* L.) in Konkan region. M. Sc. (Ag.) Thesis, Dr. B.S. Konkan Krishi Vidyapeeth, Dapoli, Ratnagiri (M.S.). 2008.
7. Morgan ED. Azadirachtin, a scientific gold mine. Bioorg. Med. Chem. 2009;17:4096-4105.
8. Neergaard P. Danish spp. of Alternaria and Stemphylium taxonomy, parasitism, economical significance. Oxford University Press, London/Oxford. 1945, 55.
9. Rotem J. The genus Alternaria: Biology, Epidemiology and Pathogenicity. The American Phytopathological Society. USA. 1994, 325.
10. Surviliene E, Dambrauskiene E. Effect of different active ingredients of fungicides on Alternaria spp. grown *in vitro*. Agron. Research. 2006;4:403-406.