



ISSN (E): 2277- 7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2022; 11(3): 1343-1346
© 2022 TPI

www.thepharmajournal.com

Received: 06-01-2022

Accepted: 09-02-2022

Akshaya M

Department of Genetics and
Plant Breeding, University of
Agricultural Sciences, Dharwad,
Karnataka, India

Shantakumar G

Professor, Department of
Genetics and Plant breeding
College of Agriculture,
Hanumanamatti University of
Agricultural Sciences, Dharwad,
Karnataka, India

O Sridevi

Professor, Department of
Genetics and Plant Breeding,
University of Agricultural
Sciences, Dharwad, Karnataka,
India

Sangamesh Nevani

Department of Genetics and
Plant Breeding, University of
Agricultural Sciences, Dharwad,
Karnataka, India

Corresponding Author:

Akshaya M

Department of Genetics and
Plant Breeding, University of
Agricultural Sciences, Dharwad,
Karnataka, India

Correlation and path coefficient analysis for determining interrelationships among grain yield and yield related characters in maize hybrids (*Zea mays* L.)

Akshaya M, Shantakumar G, O Sridevi and Sangamesh Nevani

DOI: <https://doi.org/10.22271/tpi.2022.v11.i3r.11477>

Abstract

The present investigation for studying correlations and path analysis estimates for grain yield and yield contributing characters in maize was carried out at Maize Research Centre and Seed Farm, Devihosur and analysis was carried out for eleven characters. Days to 50 per cent tasseling was significantly and positively correlated with days to 50 per cent silking and negatively correlated with plant height, ear height, cob length, shelling percentage, hundred grain weight and grain yield. Days to 50 per cent silking was significant and negatively correlated with plant height, ear height, shelling percentage and grain yield. Hundred grain weight exhibited significant positive association with plant height, ear height, cob length, cob girth, number of kernel rows per ear, number of kernels per row and grain yield. The character grain yield showed significant and negative correlation with days to 50 per cent tasseling, days to 50 per cent silking and positive correlation with plant height, ear height, cob length, cob girth, number of kernel rows per cob, number of kernels per row and hundred grain weight. Path analysis revealed that direct positive effects were shown by number of kernels per ear followed by cob length, days to fifty per cent tasseling, ear height and cob girth. Negative direct effects were shown by days to fifty per cent silking, days to 75 per cent dry husk, plant height, number of kernels per row and shelling percentage.

Keywords: Maize, correlation, path analysis

1. Introduction

Maize (*Zea mays* L., $2n=20$) is the world's leading crop and is widely cultivated as cereal grain that was domesticated in Central America. Globally, maize is known as queen of cereals because of its highest genetic yield potential, Maize is the only food crop that can be grown in diverse seasons and ecologies. Maize has become a staple food to many parts of the world with total production of maize surpassing that of wheat or rice in addition to being consumed directly by humans. Maize is also used for corn ethanol, animal feed and other maize products, such as corn starch and corn syrup. Determination of correlation and path coefficients between yield and yield traits is important for the selection of favorable plant types for effective maize breeding programs. Correlation coefficients show associations among independent characteristics and the degree of linear relation between these characteristics. It is not sufficient to describe this relationship when the causal association among characteristics is needed. Path analysis is used to determine the amount of direct and indirect effect of the causal components on the effect component. Keeping this in view, the present study was undertaken to study the extent of relationships among yield and its components for enhancing the usefulness of selection.

2. Materials and Methods

The present investigation for studying correlations and path analysis estimates for grain yield and yield contributing characters in maize was carried out during *kharif* 2020, summer 2021 at Maize Research Centre and Seed Farm, Devihosur. Phenotypic and genotypic correlations were worked out in 23 parental inbred lines and the pooled analysis over was carried out and presented in Tables 1 and 2 respectively. In general, phenotypic correlations were of higher magnitude than the corresponding phenotypic values and hence only the genotypic correlations are discussed hereunder. Path coefficient analysis allows separation of the direct effect and their indirect effects through other attributes by partitioning the correlations (Wright, 1921)^[13]. Hence, the path coefficient analysis was undertaken to know the direct and indirect effects in maize.

3. Results and Discussion

Grafius (1959) ^[1] revealed that there may not be genes controlling for yield as such but exhibit only through its components. Hence correlation analysis provides the reliable information on nature and magnitude of the association of different component characters with grain yield, which is regarded as highly complex trait in which the breeder is ultimately interested. So, there is scope for the plant breeders to identify which of the characters are correlated with yield and also the relationship among themselves, so that breeders can ultimately increase yield by selecting these component traits.

Association of days to 50 per cent tasseling was highly significant positive correlation with days to 50 per cent silking (0.601) and significant negative association for days to 50 per cent tasseling was observed with plant height (-0.160), ear height (-0.340), cob length (-0.124), shelling percentage (-0.240), hundred grain weight (-0.123) and grain yield (-0.225) at phenotypic level. Similar results for association of grain yield per plant with days to 50 percent tasseling were also reported by Raghu *et al.* (2011) ^[7], Ravi *et al.* (2012) ^[8] and Sharma *et al.* (2014) ^[10]. Days to 50 per cent tasseling in pooled analysis showed highly significant positive correlation with days to 50 per cent silking (0.914) and number of kernels per row (0.129). Significant negative association for days to 50 per cent tasseling was observed with plant height (-0.319), ear height (-0.514), cob length (-0.188), Cob girth (-0.158), shelling percentage (-0.597), hundred grain weight (-0.134) and grain yield (-0.254).

Pooled analysis revealed that association of days to 50 per cent tasseling was significant positive correlation with number of kernels per row (0.140) and significant negative association for days to 50 per cent silking was observed with plant height (-0.215), ear height (-0.356), shelling percentage (-0.261) and grain yield (-0.187). Whereas days to 75 per cent dry husk (0.014) and number of kernels per cob (0.056) had a non-significant positive association with this trait. Such finding was also observed by Reddy and Jabeen (2016) ^[9] and Jodage *et al.* (2017) ^[2]. days to 50 per cent silking was a significant positive correlation with number of kernels per row (0.195) and a significant negative association for days to 50 per cent silking was observed with days to 75 per cent dry husk (-0.145), plant height (-0.338), ear height (-0.493), cob length (-0.152), cob girth (-0.152), shelling percentage (-0.436), hundred-grain weight (-0.120) and grain yield.

The association of number of kernel rows per cob was significant and positive with plant height, ear height, cob length, cob girth, number of kernels per cob, number of kernels per row, hundred-grain weight and grain yield in the pooled analysis. Hundred seed weight exhibited significant positive association with plant height, ear height, cob length, cob girth, number of kernel rows per cob, number of kernels per row, shelling percentage and grain yield. The character grain yield per plant showed significant and negative correlation with days to 50 per cent tasseling, days to 50 per cent silking and positive correlation with plant height, ear height, cob length, cob girth, number of kernel rows per cob,

number of kernels per row and hundred-grain weight in the pooled analysis. Similar results were reported by Nataraj *et al.* (2014) ^[6] and Sharma *et al.* (2014) ^[10].

Path analysis revealed that direct positive effects were shown by number of kernels per ear (2.000), cob length (1.702), days to fifty per cent tasseling (0.930), ear height (0.100) and cob girth (0.096). Negative direct effects were shown by days to fifty per cent silking, days to 75 per cent dry husk, plant height, number of kernels per row and shelling percentage. Positive indirect effect of days to fifty per cent tasseling on grain yield was observed via days to fifty per cent silking, days to 75 per cent dry husk and number of kernels per row. A highly negative indirect effect was shown by shelling percentage (-0.555) followed by ear height (-0.478) and plant height (-0.297). Days to fifty per cent silking showed highly positive indirect on grain yield through shelling percentage (0.908) followed by plant height (0.705). It was negative for the characters *viz.*, days to fifty per cent tasseling (-1.903), number of kernels per row (-0.405) and number of kernels per ear (-0.183).

Days to 75 per cent dry husk showed a high positive indirect effect on grain yield via days to fifty per cent silking and a high negative indirect effect through number of kernels per ear. Ear height showed a positive indirect effect on grain yield through all most all the characters except days to fifty per cent tasseling and days to fifty per cent silking. Cob length showed a highly positive indirect effect on grain yield via ear height (1.122) followed by number of kernels per ear. Negative indirect effect was shown by days to fifty per cent tasseling and days to fifty per cent silking. Cob girth showed the highest positive indirect effect through shelling percentage (0.070) followed by number of kernels per ear (0.050). Number of kernels per ear showed positive indirect effect on grain yield with all the characters studied except days to fifty per cent tasseling. Number of kernels per row showed negative indirect effect through all the traits under study except shelling percentage (0.012). Shelling percentage showed positive indirect effect on grain yield through days to fifty per cent tasseling (1.092), days to fifty per cent silking (0.798) and number of kernels per row (0.027). Highest negative indirect effect was shown by cob girth (-1.327) followed by cob length (-0.845).

The present investigation clearly revealed that the characters number of kernels per ear, cob length and cob girth had a strong association with grain weight and positive highest direct and an indirect effect through other component traits. Vaezi *et al.* (2000) ^[12] reported that the bulk of indirect effect on grain yield understudy was exerted by the traits like ear length and ear girth. This indicated direct selection for these characters will enhance the breeding efficiency for grain yield in maize. Mohamadi *et al.* (2003); Kumar *et al.* (1999) reported that number of kernels per ear had a maximum positive direct effect on yield followed by cob length, plant height, cob girth and days to fifty per cent silking. Sofi and Rather (2006) ^[11], Najeeb *et al.* (2009) ^[5] also reported the importance of these characters in path analysis.

Table 1: Phenotypic (P) and Genotypic (G) pooled correlations for 12 characters in maize

Source		Days to 50% Tasseling	Days to 50% silking	Days to maturity	Plant height (cm)	Ear height (cm)	Ear length (cm)	Ear girth (cm)	Number of kernel rows per ear	Number of kernels per row	Shelling percentage (%)	100-grain weight (g)	Grain yield (kg/ha)
Days to 50% tasseling	P	1.00	0.601**	0.015	-0.160*	-0.340*	-0.124*	-0.070	-0.045	0.063	-0.240*	-0.123*	-0.225*
	G	1.00	0.914**	0.079	-0.319*	-0.514*	-0.188*	0.158*	-0.030	0.129*	-0.597*	-0.134*	-0.254*
Days to 50% silking	P		1.00	0.014	-0.215*	-0.356*	-0.094	-0.072	0.056	0.140*	-0.261*	-0.084	-0.187*
	G		1.00	-0.145*	-0.338*	-0.493*	-0.152*	0.152*	0.088	0.195*	-0.436*	-0.120*	-0.245*
Days to maturity	P			1.00	0.076	0.120	0.138*	0.078	0.111*	0.064	0.080	0.111*	0.106*
	G			1.00	0.070	0.336*	0.145	-0.048	0.366*	0.100*	0.012	0.037	0.003
Plant height (cm)	P				1.00	0.662**	0.462*	0.258*	0.283*	0.266*	0.142*	0.321*	0.454*
	G				1.00	0.758**	0.580*	0.336*	0.382*	0.334*	0.243*	0.400*	0.580*
Ear height (cm)	P					1.00	0.499*	0.241*	0.333*	0.334*	0.193*	0.287*	0.482*
	G					1.00	0.659**	0.289*	0.387*	0.439*	0.346*	0.367*	0.550*
cob length (cm)	P						1.00	0.184*	0.173*	0.447*	0.161*	0.349*	0.544*
	G						1.00	0.242*	0.276*	0.580*	0.462*	0.419*	0.627*
cob girth (cm)	P							1.00	0.381*	0.011	0.329*	0.429*	0.443*
	G							1.00	0.518*	0.002	0.725**	0.558*	0.514*
Number of kernels per ear	P								1.00	0.225*	0.240*	0.459*	0.569*
	G								1.00	0.363*	0.522*	0.631*	0.690*
Number of kernels per row	P									1.00	-0.088	0.313*	0.413*
	G									1.00	-0.010	0.358*	0.452*
Shelling percentage	P											0.319*	0.394*
	G											0.733**	0.748**
100 seed weight (g)	P											1.00	0.765**
	G											1.00	0.829**

Table 2: Phenotypic (P) and Genotypic (G) pooled path for 12 characters in maize

Source		Days to 50% tasseling	Days to 50% silking	Days to maturity	Plant Height(cm)	Ear height (cm)	Cob length (cm)	cob girth (cm)	Number of kernel rows per ear	Number of kernels per row	Shelling percentage (%)	100-seed weight (g)	Grain yield (kg/ha)
Days to 50% tasseling	P	-0.046	-0.028	-0.001	0.007	0.016	0.006	0.003	0.002	-0.003	0.011	0.006	-0.226
	G	0.930	0.851	0.074	-0.297	-0.479	-0.176	-0.148	-0.029	0.120	-0.556	-0.126	-0.254
Days to 50% silking	P	-0.051	-0.084	-0.001	0.018	0.030	0.008	0.006	-0.005	-0.012	0.022	0.007	-0.188
	G	-1.903	-2.081	0.303	0.705	1.028	0.317	0.318	-0.184	-0.406	0.909	0.251	-0.245
Days to maturity	P	0.000	0.000	-0.026	-0.002	-0.003	-0.004	-0.002	-0.003	-0.002	-0.002	-0.003	0.107
	G	-0.097	0.178	-1.224	-0.086	-0.412	-0.178	0.059	-0.448	-0.122	-0.016	-0.045	0.004
Plant height (cm)	P	-0.007	-0.009	0.003	0.041	0.027	0.019	0.011	0.012	0.011	0.006	0.013	0.455
	G	0.385	0.408	-0.084	-1.206	-0.914	-0.700	-0.406	-0.461	-0.403	-0.293	-0.482	0.580
Ear height (cm)	P	-0.005	-0.005	0.002	0.010	0.015	0.008	0.004	0.005	0.005	0.003	0.004	0.483
	G	-0.052	-0.050	0.034	0.076	0.100	0.066	0.029	0.039	0.044	0.035	0.037	0.551
cob length (cm)	P	-0.028	-0.021	0.031	0.103	0.111	0.223	0.041	0.039	0.100	0.036	0.078	0.545
	G	-0.322	-0.259	0.248	0.988	1.122	1.703	0.414	0.471	0.988	0.788	0.714	0.628
cob girth (cm)	P	-0.004	-0.004	0.005	0.016	0.015	0.011	0.061	0.023	0.001	0.020	0.026	0.444
	G	-0.015	-0.015	-0.005	0.033	0.028	0.023	0.097	0.050	0.000	0.070	0.054	0.515
Number of kernel rows per ear	P	-0.011	0.013	0.026	0.066	0.078	0.041	0.089	0.233	0.053	0.056	0.107	0.569
	G	-0.062	0.177	0.733	0.765	0.775	0.554	1.038	2.001	0.727	1.046	1.263	0.691
Number of kernels per row	P	0.008	0.018	0.008	0.034	0.043	0.058	0.002	0.029	0.129	-0.011	0.040	0.413
	G	-0.105	-0.158	-0.081	-0.270	-0.355	-0.469	-0.002	-0.294	-0.809	0.012	-0.290	0.452
Shelling percentage	P	-0.027	-0.029	0.009	0.016	0.021	0.018	0.036	0.026	-0.010	0.110	0.035	0.395
	G	1.092	0.799	-0.024	-0.445	-0.634	-0.846	-1.327	-0.956	0.028	-1.829	-1.341	0.748
100 seed weight (g)	P	-0.056	-0.038	0.051	0.145	0.130	0.158	0.194	0.208	0.141	0.144	0.451	0.766
	G	-0.107	-0.096	0.029	0.318	0.292	0.333	0.443	0.501	0.284	0.582	0.794	0.829

4. Conclusion

The results obtained in this investigation revealed the occurrence of considerable positive as well as negative direct and indirect effects by various characters on the yield of Maize through one or other characters. Thus, it can be concluded that the characters mentioned above should be duly considered at the time of formulation of selection strategy to develop high yielding hybrids in Maize.

5. Acknowledgements

The authors' wish to express their gratitude to Dr. G Shantakumar Eminent Maize breeder and My Batchmates Anil, Maruthi, Srivijay, Ashritha, Supriya, Sharada and my dear juniors.

6. References

1. Grafius JE. Heterosis in barely. Journal of Agronomy. 1959;51:551-554.
2. Jodage K, Kuchanur PH, Zaidi PH, Patil A, Seetharam K, Vinayan MT. Association and path analysis for grain yield and its attributing traits under heat stress condition in tropical maize (*Zea mays* L.). Electronic journal of plant breeding. 2017;8(1):336-341.
3. Kumar A, Gangashetti MG, Dahiya A. Analysis of direct and indirect effects for quantitative traits in diallel crosses of maize (*Zea mays*). Annuals of Biology. 1997;15(2):173-176.
4. Mohammadi SA, Prasanna BM Singh NN. Sequential path model for determining interrelationships among grain yield and related characters in maize. Crop Science. 2003;43(5):1690-1697.
5. Najeeb, Sofi, Rather AG, Parray GA, Sheikh FA, Razvi SM. Studies on genetic variability, genotypic correlation and path coefficient analysis in maize under the high-altitude temperate conditions of Kashmir. Maize Genet. Coop. News Lett., 2009, 83.
6. Nataraj V, Shahi JP Agarwal V. Correlation and path analysis in certain inbred genotypes of maize (*Zea Mays* L.) at Varanasi. International journal of innovative research. 2007;3(1):14-17.
7. Raghu B, Suresh J, Kumar SS, Saidaiah P. Character association and path analysis in maize (*Zea mays* L.). Madras agricultural journal 98(1-3), 7-9.
8. Ravi VM, Chikkalingaiah Shailaja, H. Correlation study for protein content, grain yield and yield contributing traits in quality protein maize (*Zea mays* L.). Electronic journal of plant Breeding. 2012;3(1):649-651.
9. Reddy VR, Jabeen F. Narrow sense heritability, correlation and path analysis in maize (*Zea mays* L.). Sabrao journal of breeding. 2016;48(2):120-126.
10. Sharma T, Kumar A, Dwivedi SC, Vyas RP. Estimate of genetic factors and correlation analysis in maize (*Zea mays* L.). Plant Archeives. 2014;14(1):19-21.
11. Sofi P, Rather AG. Genetic analysis of yield traits in local and CIMMYT inbred line crosses using linextester analysis in maize (*Zea mays* L.). Asian Journal of Plant science 2006.
12. Vaezi SH, Abd-Mishani C, Yazdi-Samadi B Ghannadha MR. Correlation and path analysis of grain yield and its components in maize. Iranian Journal of agricultural science. 2000;31(1):71-83.
13. Wright S. Systems of mating and The biometric relations between parent and offspring. Genetics. 1921;6(2):111.