



ISSN (E): 2277-7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2022; SP-11(5): 1479-1483
© 2022 TPI
www.thepharmajournal.com
Received: 15-03-2022
Accepted: 21-04-2022

Ushamalini C

Associate Professor
(Plant Pathology), Forest
College and Research Institute,
Mettupalayam, Tamil Nadu,
India

Dr. Meena B

Professor (Plant Pathology),
Coconut Research Station,
Aliyar Nagar, Tamil Nadu, India

Sudhalakshmi R

Assistant Professor (Soil Science
and Agricultural Chemistry),
Coconut Research Station,
Aliyar Naga, Tamil Nadu, India

Dr. Indra N

Assistant Professor
(Plant Pathology), Forest
College and Research Institute,
Mettupalayam, Tamil Nadu,
India

Development of IPM module for the management of soil borne diseases in groundnut

Ushamalini C, Dr. Meena B, Sudhalakshmi R and Dr. Indra N

Abstract

Ground nut (*Arachis hypogea* L.), is one of the most important oil seed crop of India, is affected by several soil borne diseases viz., root rot and stem rot. A study was conducted for the development of IPM module for the management of these soil borne fungicides using fungicides and biocontrol agents. Among the nine modules tested, module with deep summer ploughing using mould board plough combined with soil application of *Trichoderma* @ 4 kg/ ha enriched in 250 kg FYM/ha and seed treatment with Tebuconazole 2DS recorded minimum incidence of stem rot (9.5%) with an pod and haulm yield of 2566 kg/ha and 6428 kg /ha respectively. This treatment also recorded minimum incidence of late leaf spot (Grade 4) and rust (Grade 5).

Keywords: IPM module, soil borne, *Arachis hypogea* L.

Introduction

Ground nut (*Arachis hypogea* L.), is one of the most important oil seed crop of India, contributing about 36% of the world oil seed production. The major peanut growing states area Gujarat, Andra Pradesh, Tamil Nadu, Rajasthan and Maharashtra which contributes around 90% of the area and production. Seed and soil borne diseases causes severe seedling mortality resulting in the patchy crop stand mostly in the sandy loam soils and reduces yield from 25 to 50%. Among the soil borne diseases, collar rot (*Aspergillus niger* Van Tieghem), stem rot (*Sclerotium rolfsii* Sacc) and dry root rot (*Macrophomina phaseolina* (Tassi Goid) = *Rhizoctonia bataticola* (Taub.) Butler) have been recognized as a major diseases. Among the foliar diseases ELS, LLS and rust are commercially important and cause yield losses up to 50 per cent. The fungicides viz., azoxystrobin, chlorothalonil and tebuconazole were found to be effective for the management of leaf spot and stem rot in groundnut (Hagan *et al.*, 2010) [5]. Wann *et al.*, 2011 [8] reported the management of leaf spot in ground nut using fungicides. Integrated management of diseases by combining chemical and biological method would be an effective method for the management of diseases compared to adopting single method. With this view, the present work was carried out to find out effective IDM module for the management stem rot and foliar diseases in ground nut.

Materials and Methods

Three field trials were conducted to test the efficacy of different fungicides for the management of foliar diseases in groundnut during 2016-17, 2017-18 and 2018-19 in Kharif season at Coconut Research station, Aliyarnagar, TamilNadu Agricultural University (Longitude: 77.49°E latitude: 10°N MSL- 260 msl). The ground nut var CO-2, was used for sowing at 30x 10 cm spacing in 15 m² (5m x 3m) plots and the plots were laid out in Randomized Block Design (RBD) with three replications. Treatments were given as per schedule. The incidence of stem rot was recorded on 45-50 DAS by Percent Diseases Incidence (PDI). Foliar disease incidences were recorded periodically by adopting 1-9 scale (Subramanyam *et al.*, 1995) [7]. The pod and haulm yield were also recorded after harvest.

Corresponding Author

Ushamalini C

Associate Professor
(Plant Pathology), Forest
College and Research Institute,
Mettupalayam, Tamil Nadu,
India

The treatments were

T ₁	(M5): Popular variety + Deep summer ploughing with mold board plough + Soil application of <i>Trichoderma viride</i> @ 4 kg/ ha enriched in FYM+Seed treatment with Tebuconazole @ 1.5 g/kg of seed
T ₂	(M10): Popular Variety + Deep summer ploughing with mold board plough + Soil application of <i>T. viride</i> @ 4 kg/ ha enriched in FYM + Seed treatment with <i>T. viride</i> @ 10 kg of seed + Soil application of Trichoderma @ 4 kg/ ha enriched in FYM at 35 days after sowing (DAS)
T ₃	(M11): Popular Variety + Deep summer ploughing with mold board plough + Soil application of <i>T. viride</i> @ 4 kg/ ha enriched in FYM + Seed treatment with Tebuconazole @ 1.5 g/ kg of seed + Soil application of Trichoderma @ 4 kg/ ha enriched in FYM at 35 days after sowing (DAS)
T ₄	Deep summer ploughing with mold board plough + Soil application of <i>T. viride</i> @ 4 kg/ ha enriched in FYM + Seed treatment with Tebuconazole 2DS @ 1.5 g/ kg of seeds followed by seed treatment with PGPR @ 625 g for per ha of seeds + Soil application of <i>T. viride</i> @ 4 kg/ ha enriched in 250 kg FYM/ ha at 35 and 70 DAS
T ₅	(M15) : Popular Variety + Deep summer ploughing with mold board plough + Soil application of <i>T. viride</i> @ 4 kg/ ha enriched in FYM + Seed treatment with PGPR @ 625g/ for per ha of seeds + Soil application of <i>T. viride</i> @ 4 kg/ ha enriched in FYM at 35 days after sowing (DAS) + Soil application of <i>T. viride</i> @ 4 kg/ ha enriched in FYM at 80 days after sowing (DAS)
T ₆	(M17) : Popular Variety + Deep summer ploughing with mold board plough + Soil application of <i>T. viride</i> @ 4 kg/ ha enriched in FYM + Seed treatment with Tebuconazole @ 1.5 g/kg of seeds + Soil application of <i>T. viride</i> @ 4 kg/ ha enriched in FYM at 35 days after sowing (DAS) + Soil application of <i>T. viridea</i> @ 4 kg/ ha enriched in FYM at 80 days after sowing (DAS)
T ₇	Deep summer ploughing
T ₈	Farmers Practice (respective centre)
T ₉	Control (Without any treatments)

Results

The results of the trial conducted during 2017-18 indicated that, among the nine treatments tested, T₄ treatment *i.e.* Deep summer ploughing with mold board plough+ Soil application of *Trichoderma* @ 4 kg/ ha enriched in 250 kg FYM/ha+ Seed treatment with Tebuconazole 2DS recorded minimum incidence of stem rot (8.47%) with an pod and haulm yield of 3099 kg/ha and 5997 kg /ha respectively. This treatment also recorded minimum incidence of Late leaf spot (Grade 4) and rust (Grade 6). This treatment is followed by T₅ treatment *i.e.* Popular Variety + Deep summer ploughing with mold board

plough + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM + Seed treatment with PGPR @ 625g/ for per ha of seeds + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM at 35 days after sowing (DAS) + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM at 80 days after sowing (DAS) that recorded an incidence of 17.7% and pod and haulm yield of 3057 kg/ha and 5989 kg/ha respectively, while in control the stem rot incidence was 36.7%.with pod and haulm yield of 2062 and 5308 kg/ha respectively (Table 1). In control plots, the incidence of late leaf spot and rust were maximum (Grade 8).

Table 1: Validation of Management module for the soil borne diseases (2017-18)

Treatments	Germination (%)	Collar rot (%)	Stem rot (%)				Dry root rot (%)				Foliar diseases (1-9 scale)		Pod rot (%)	Pod yield (kg/ha)	Haulm yield (kg/ha)
			20-30 DAS	40-45 DAS	70-75 DAS	At harvest	30-35 DAS	60-65 DAS	85-90 DAS	At harvest	LLS	Rust			
T ₁	85.1	0	0	0	19.7 (26.4)	0	0	0	0	4	7	0	3003	5943	
T ₂	90.2	0	0	0	23.3 (28.8)	0	0	0	0	5	7	0	3016	5729	
T ₃	87.5	0	0	0	21.3 (27.4)	0	0	0	0	5	7	0	3055	5737	
T ₄	84.9	0	0	0	8.47 (10.8)	0	0	0	0	4	6	0	3099	5997	
T ₅	86.8	0	0	0	17.7 (24.8)	0	0	0	0	4	6	0	3057	5989	
T ₆	91.2	0	0	0	25.6 (30.4)	0	0	0	0	4	7	0	3018	5670	
T ₇	89.3	0	0	0	28.7 (32.3)	0	0	0	0	4	6	0	2983	5676	
T ₈	88.7	0	0	0	18.9 (25.8)	0	0	0	0	5	7	0	2495	5651	
T ₉	89.1	0	0	0	36.7 (37.2)	0	0	0	0	8	8	0	2062	5308	
SEm+1					1.80								66.9	374.8	
CD at 5%					3.79								140.6	787.6	
CV%					8.09								2.86	7.99	

The results of the trial conducted during 2018-19 indicated that, among the nine treatments tested, T₄ treatment *i.e.* : Deep summer ploughing with mold board plough+ Soil application of *Trichoderma* @ 4 kg/ ha enriched in 250 kg FYM/ha+ Seed treatment with Tebuconazole 2DS recorded minimum

incidence of stem rot (12.5%) with an pod and haulm yield of 2342 kg/ha and 6650 kg /ha respectively. This treatment also recorded minimum incidence of Late leaf spot (Grade 3) and rust (Grade 4). This treatment is followed by T₅ treatment *i.e.* Popular Variety + Deep summer ploughing with mold board

plough + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM + Seed treatment with PGPR @ 625g/ for per ha of seeds + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM at 35 days after sowing (DAS) + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM at 80 days after sowing (DAS) that recorded an incidence of 13.6% and pod and

haulm yield of 2286 kg/ha and 6180 kg/ha respectively, while in control the stem rot incidence was 30.6%.with pod and haulm yield of 1231 and 4398 kg/ha respectively. In control plots, the incidence of late leaf spot and rust were Grade 7 and Grade 6 respectively (Table 2).

Table 2: Validation of management modules for soil borne diseases (2018-19)

Treatments	Germination (%)	Collar rot (%)	Stem rot (%)				Dry root rot (%)				Foliar diseases (1-9 scale)		Pod rot (%)	Pod yield (kg/ha)	Haulm yield (kg/ha)
			20-30 DAS	40-45 DAS	70-75 DAS	At harvest	30-35 DAS	60-65 DAS	85-90 DAS	At harvest	LLS	Rust			
T ₁	85.7	0	0	0	19.6 (26.33)	0	0	0	0	4	5	0	2084	5980	
T ₂	83.7	0	0	0	16.9 (24.33)	0	0	0	0	3	5	0	2190	6092	
T ₃	84.4	0	0	0	15.3 (23.00)	0	0	0	0	4	6	0	2262	6149	
T ₄	86.9	0	0	0	12.5 (20.77)	0	0	0	0	3	4	0	2342	6650	
T ₅	85.5	0	0	0	13.6 (21.64)	0	0	0	0	4	5	0	2286	6180	
T ₆	84.1	0	0	0	20.1 (26.64)	0	0	0	0	4	4	0	2042	5720	
T ₇	81.2	0	0	0	23.5 (29.00)	0	0	0	0	5	6	0	1582	4872	
T ₈	85.5	0	0	0	22.4 (28.25)	0	0	0	0	3	5	0	1849	5290	
T ₉	79.8	0	0	0	30.6 (33.59)	0	0	0	0	7	6	0	1231	4398	
SEm±1	1.05				1.90								136.4	682.94	
CD at 5%	2.22				4.04								264.33	1553.43	
CV%	1.53				12.01								7.43	16.09	

The results of the trial conducted during 2019-20 indicated that, among the nine treatments tested, T₄ treatment *i.e.* Deep summer ploughing with mold board plough+ Soil application of *Trichoderma* @ 4 kg/ ha enriched in 250 kg FYM/ha+ Seed treatment with Tebuconazole 2DS recorded minimum incidence of stem rot (7.5%) with an pod and haulm yield of 2226 kg/ha and 6637 kg /ha respectively. This treatment also recorded minimum incidence of Late leaf spot (Grade 4) and rust (Grade 6). This treatment is followed by T₅ treatment *i.e.* Popular Variety + Deep summer ploughing with mold board plough + Soil application of *T. viride* @ 4 kg/ ha enriched in

FYM + Seed treatment with PGPR @ 625g/ for per ha of seeds + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM at 35 days after sowing (DAS) + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM at 80 days after sowing (DAS) that recorded an incidence of 15.5% and pod and haulm yield of 2003 kg/ha and 6594 kg/ha respectively, while in control the stem rot incidence was 37.1%.with pod and haulm yield of 1158 and 4436 kg/ha respectively (Table 3). In control plots, the incidence of late leaf spot and rust were maximum (Grade 8).

Table 3: Validation of management modules for soil borne diseases (2019-20)

Treatments	Germination (%)	Collar rot (%)	Stem rot (%)				Dry root rot (%)				Foliar diseases (1-9 scale)		Pod rot (%)	Pod yield (kg/ha)	Haulm yield (kg/ha)
			20-30 DAS	40-45 DAS	70-75 DAS	At harvest	30-35 DAS	60-65 DAS	85-90 DAS	At harvest	LLS	Rust			
T ₁	92.1	0	0	0	20.0 (26.5)	0	0	0	0	4	7	0	1923	6076	
T ₂	90.2	0	0	0	22.7 (28.3)	0	0	0	0	3	5	0	1832	6306	
T ₃	85.5	0	0	0	21.7 (27.7)	0	0	0	0	5	7	0	1821	6381	
T ₄	87.9	0	0	0	7.5 (15.9)	0	0	0	0	4	6	0	2226	6637	
T ₅	88.8	0	0	0	15.5 (23.1)	0	0	0	0	4	5	0	2003	6594	
T ₆	87.2	0	0	0	21.5 (27.5)	0	0	0	0	4	7	0	1842	6217	
T ₇	86.3	0	0	0	26.7 (31.1)	0	0	0	0	4	6	0	1352	5459	

T ₈	90.2	0	0	0	22.8 (28.4)	0	0	0	0	5	7	0	1494	5259
T ₉	89.6	0	0	0	37.1 (37.5)	0	0	0	0	8	8	0	1158	4436
SEm+1					2.05								129.50	779.94
CD at 5%					4.36								274.53	1653.43
CV%					9.22								7.59	16.09

The pooled results of the three years (2017-18, 2018-19 and 2019-20) trial data indicated that, all the treatments recorded germination percentage of more than 80.0 per cent. Among the nine treatments tested, T₄ treatment i.e. Deep summer ploughing with mould board plough + Soil application of *Trichoderma* @ 4 kg/ ha enriched in 250 kg FYM/ha+ Seed treatment with Tebuconazole 2DS recorded minimum incidence of stem rot (9.5%) with an pod and haulm yield of 2566 kg/ha and 6428 kg /ha respectively. This treatment also recorded minimum incidence of Late leaf spot (Grade 4) and rust (Grade 5). This treatment is followed by T₅ treatment i.e. Popular Variety + Deep summer ploughing with mold board

plough + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM + Seed treatment with PGPR @ 625g/ for per ha of seeds + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM at 35 days after sowing (DAS) + Soil application of *T. viride* @ 4 kg/ ha enriched in FYM at 80 days after sowing (DAS) that recorded an incidence of 15.6% and pod and haulm yield of 2499 kg/ha and 6254 kg/ha respectively, while in control the stem rot incidence was 34.8%.with pod and haulm yield of 1484 and 4714 kg/ha respectively. In control plots, the incidence of late leaf spot and rust were Grade 8 and grade 7 respectively (Table 4).

Table 4: Validation of management modules for soil borne diseases (Pooled data) Centre: CRS, Aliyarnagar

Treatments	Germination (%)	Collar rot (%)	Stem rot (%)				Dry root rot (%)				Foliar diseases (1-9 scale)		Pod rot (%)	Pod yield (kg/ha)	Haulm yield (kg/ha)
			20-30 DAS	40-45 DAS	70-75 DAS	At harvest	30-35 DAS	60-65 DAS	85-90 DAS	At harvest	LLS	Rust			
T ₁	85.7	0	0	0	0	19.8	0	0	0	0	4	6	0	2337	6000
T ₂	83.7	0	0	0	0	21.0	0	0	0	0	4	6	0	2346	6042
T ₃	84.4	0	0	0	0	19.4	0	0	0	0	5	7	0	2379	6089
T ₄	86.9	0	0	0	0	9.5	0	0	0	0	4	5	0	2556	6428
T ₅	85.5	0	0	0	0	15.6	0	0	0	0	4	5	0	2449	6254
T ₆	84.1	0	0	0	0	22.4	0	0	0	0	4	6	0	2301	5869
T ₇	81.2	0	0	0	0	26.3	0	0	0	0	4	6	0	1972	5336
T ₈	85.5	0	0	0	0	21.4	0	0	0	0	4	6	0	1946	5400
T ₉	79.8	0	0	0	0	34.8	0	0	0	0	8	7	0	1484	4714
SEm+1						1.9								111.9	577.4
CD at 5%						4.1								226.5	1331.5
CV%						9.8								6.1	13.4

The pooled result of trials indicated that combing of cultural, chemical and biological method i.e. Deep summer ploughing with mould board plough + Soil application of *Trichoderma* @ 4 kg/ ha enriched in 250 kg FYM/ha+ Seed treatment with Tebuconazole 2DS is highly effective in reducing the incidence of stem rot and also foliar diseases.

The effect of seed treatment in increasing the yield was reported by several workers. Dandnaik *et al.*, 2009^[1] reported that seed treatment with hexaconazole gave higher pod yield in ground nut. De *et al.*, (2003)^[2] with carbendazim + Thiram in linseed against wilt, Ramkishore and Singh (2008) in linseed against wilt, Dudi and Lodha (2003)^[3] in pea nut against seedling diseases, Dutta and Das (2002)^[4] against collar rot in tomato were also already reported. Integrated disease management of foliar and soil borne diseases with fungicides, Castor cake and *Trichoderma* in ground nut was reported by Jadon *et al.*, 2017^[6]. They reported that, lowest incidence of soil borne diseases was recorded in seed treatment with mancozeb and seed treatment with tebuconazole compared to untreated control.

References

- Dandnaik BP, Patil DT, Chavanand MH, Dandnaik AB. Chemical control of stem rot of groundnut caused by *Sclerotium rolfsii*. Journal of Mycology and Plant Pathology. 2009;39:185.
- De RK, Dwivedi RP, Udit Narain. Biological control of lentil wilt caused by *Fusarium oxysporum* fsp. lentis. Annals of Plant Protection Sciences. 2003;11:46-52.
- Dudi J, Lodha PC. Compatibility of seed dressing fungicides with rhizobium in groundnut. National seminar: Stress management in oil seeds, 2003, Jan 28-30, 24-25.
- Dutta P, Das BC. Management of collar rot of tomato by *Trichoderma* spp. And chemicals. Indian Phytopathology 2002;55:235-237.
- Hagan AK, Campbell HL, Bowen KL, Wells L, Goodman R. Managing early leaf spot and stem rot with reduced fungicide inputs on disease resistant peanut cultivars. Peanut Science. 2010;37:129-136.
- Jadon KS, Thirumalaiswamy PP, Vinod Kumar, Koradia VG, Padavi RD. Integrated management of major foliar and soil borne diseases of pea nut (*Arachis hypogea* L.) with fungicides, *Trichoderma* and Castor cake. Int. J Curr. Microbiol. App. Sci, 2017;6(12):1884-1899.
- Subrahmanyam P, McDonald D, Waliyar E, Reddy LJ, Nigam SN, Gibbons RW, *et al.* Screening methods and sources of resistance to rust and late leaf spot of groundnut (In En. Summaries in En, Fr, Sp, and Pt.). Information Bulletin no. 47. Patancheru 502 324, Andhra

Pradesh, India: International Crops Research Institute for the Semi-Arid Tropics. Thind T S 2008, 1995, 24.

8. Wann DQ, Tubbs RS, Culbreath AK. Genotype and approved fungicides evaluation for reducing leaf spot diseases in organically-managed peanut. Plant Health Progress. (October): PHP-2010-1027-01-RS, 2011.