



ISSN (E): 2277-7695
ISSN (P): 2349-8242
NAAS Rating: 5.23
TPI 2022; SP-11(6): 327-335
© 2022 TPI
www.thepharmajournal.com
Received: 25-03-2022
Accepted: 29-04-2022

More BS
Department of Soil Science and
Agricultural Chemistry, Collage
of Agriculture, Latur,
Maharashtra, India

Adsul PB
Department of Soil Science and
Agricultural Chemistry, Collage
of Agriculture, Latur,
Maharashtra, India

Salunke PR
Department of Soil Science and
Agricultural Chemistry, Collage
of Agriculture, Latur,
Maharashtra, India

Lingayat NR
Department of Soil Science and
Agricultural Chemistry, Collage
of Agriculture, Latur,
Maharashtra, India

Mutkule US
Department of Soil Science and
Agricultural Chemistry, Collage
of Agriculture, Latur,
Maharashtra, India

Corresponding Author
More BS
Department of Soil Science and
Agricultural Chemistry, Collage
of Agriculture, Latur,
Maharashtra, India

To study the physico-chemical properties of soils of soybean growing areas from Renapur Tahsil of Latur district, Maharashtra

More BS, Adsul PB, Salunke PR, Lingayat NR and Mutkule US

Abstract

The present investigation entitled “Soil Fertility Evaluation of Soybean Growing Areas from Renapur Tahsil of Latur District” was undertaken for assessing the fertility status of soils from studied area. For this purpose, total 100 soil samples collected from 20 villages and 5 soil samples from each village were distributed according to their respective depth. Out of which 56, 26 and 18 number of samples were identified and categorized as Inceptisol, Entisol and Vertisol soil orders, respectively. These soil samples were analyzed for physico-chemical properties that is pH, electrical conductivity, organic carbon and calcium carbonate. The soils of Renapur tahsil were neutral to moderately alkaline in reaction, EC of the soil was no deleterious effect on crop. Low to moderate in organic carbon content and non-calcareous to calcareous in nature.

Keywords: PH, EC, OC, CaCO₃

Introduction

Latur is one of the districts in Marathwada region. It comes under Aurangabad administrative division. Latur district comprises total 10 tahsil, out of that Renapur tahsil selected for the study of soil fertility status. Major crops grown in this region were sugarcane, pigeon pea, gram, sorghum, wheat, soybean, groundnut, sunflower and some horticulture crops.

The physico-chemical properties like pH, EC, calcium carbonate and organic carbon are essential because they influence on availability of nutrients in soil and as a result crop growth and output. The organic carbon is the store house of all the plant nutrients. It provides good aeration, increase microbial activity, water holding capacity maintain the soil pH and calcium carbonate content in the soils. Supply of essential nutrient from the soil can be augmented by proper management of these property. For sustainable agriculture production the information on soil characterization in relation to fertility status of the soil of the region will be useful.

Materials and Methods

To evaluate soil fertility status of Renapur tahsil, 100 soil samples were collected from soybean growing area of tahsil. The twenty representative villages (Arajkheda, Sangavi, Gavhan, Andalgaon, bavchi, Itti, Nagapur, Kalewadi, Khanapur Pohergaon, Fardpur, Wala, Kamkheda, Sarola, Narwatwadi, kumbharwadi, Renapur, Shelu-kh, Kharola and harwadi) are selected and from each village 5 samples were collected. one hundred representative surface soil samples (15-20 cm) depth were collected. The soils were grouped into different orders according to USDA classification. Out of 100 samples, 56 soil samples were in Inceptisol, 26 soil samples were in Entisol and 18 soil samples were in Vertisol order, respectively.

The samples were analyzed for the physico-chemical properties of these soils, pH and Electrical conductivity (EC) in 1:2.5 soil water suspension (Jackson, 1978) [5]. The organic carbon estimated by modified method of Walkley and Black method (Jackson, 1978) [5]. Calcium carbonate determined by rapid titration method (Piper, 1966) [11].

Results and Discussion

Out of all one hundred soil samples, 56 soil samples under Inceptisol, while 26 and 18 soil samples categorized in Entisol and Vertisol, respectively. The data on physico-chemical properties of Inceptisol, Entisol and Vertisol soils are presented in table 1, 2 and 3, respectively.

Physico-chemical properties of Inceptisol soils in Renapur tahsil.

The physico-chemical properties of Inceptisol soil are presented in table 1. The lowest pH value (6.54) was recorded in soil sample (V₇S₅) from Andalgaon village, while the highest pH (8.14) of the soil sample (V₁₈S₂) was observed in Wala village. In Inceptisol soil the pH ranged from 6.54 to 8.14 with an average value 7.54. The data presented in table 4 and depicted in fig 2 reported that, out of 56 soil sample from Inceptisol, 14 samples (25%) were neutral in reaction (pH 6.5-7.5) and 42 samples (75%) were moderately alkaline in reaction (7.5-8.5).

The highest EC (1.59 dSm⁻¹) was recorded in soil sample (V₅S₁) from Arajkheda village and the lowest EC (0.27 dSm⁻¹) was observed in the soil sample (V₁₄S₁) in Gavhan. The EC was ranged from 0.27-1.59 dSm⁻¹ with a mean value of 0.65 dSm⁻¹. The data presented in table 4 and depicted in fig 3 revealed that out of 56 soil samples, 53 soil samples (94%) was no deleterious effect on crop and 3 samples (6%) were critical for germination.

Organic carbon content (Table 1) was ranged from 1.5 to 8.7 g kg⁻¹ with an average value 4.52 g kg⁻¹. The soil sample (V₁₆S₅) from Narvatwadi village showed lowest content of organic carbon (1.5 g kg⁻¹) while the highest organic carbon content (8.7 g kg⁻¹) was recorded in sample (V₅S₁) from Arajkheda village. The data presented in table 5 and depicted in fig 4 revealed that out of 56 soil samples from Inceptisol, 1 sample (2%) was very low, 23 samples (41%) were low, 23 samples (41%) were moderate, 7 samples (12%) moderately high and 2 samples (4%) were high in organic carbon content. The highest value of calcium carbonate (82 g kg⁻¹) was recorded in Faradpur village (V₁₉S₅) and lowest calcium carbonate content (20 g kg⁻¹) was observed from Khanapur village (V₁₀S₄). The calcium carbonate content in this soil were varied from 20 to 82 g kg⁻¹ with an average value of 55.13 g kg⁻¹. The data presented in table 5 and depicted in fig 4 shows that, out of 56 soil samples, 23 samples (41%) were non-calcareous (Fit for all crops) in nature and 33 samples (59%) were calcareous (Fit for all crops except citrus) in nature.

Table 1: Physico-chemical properties of Inceptisol soils in Renapur tahsil

Sr. No.	Sample No.	pH	EC (dSm ⁻¹)	OC (g kg ⁻¹)	CaCO ₃ (g kg ⁻¹)
1	V ₃ S ₁	7.04	0.48	8.2	78
2	V ₃ S ₂	7.47	0.39	3.0	74
3	V ₃ S ₃	7.46	0.63	3.8	62
4	V ₃ S ₄	7.80	1.36	4.9	69
5	V ₃ S ₅	7.55	0.69	4.1	47
6	V ₄ S ₁	7.71	1.20	6.1	39
7	V ₄ S ₂	7.78	0.69	4.9	47
8	V ₄ S ₃	7.76	0.88	4.6	46
9	V ₄ S ₄	7.56	0.52	3.9	75
10	V ₄ S ₅	7.83	0.50	3.7	50
11	V ₅ S ₁	7.96	1.59	8.7	46
12	V ₅ S ₂	7.81	0.68	4.9	75
13	V ₅ S ₄	7.58	0.36	4.0	31
14	V ₆ S ₁	7.38	0.56	4.0	29
15	V ₆ S ₂	7.50	0.59	4.5	69
16	V ₆ S ₅	7.29	0.72	3.6	59
17	V ₇ S ₁	7.62	0.71	5.2	64
18	V ₇ S ₂	7.63	0.89	3.6	64
19	V ₇ S ₄	7.56	0.73	3.4	74
20	V ₇ S ₅	6.54	0.89	2.9	43
21	V ₈ S ₁	7.48	0.46	3.4	67
22	V ₈ S ₂	7.47	0.55	4.0	53
23	V ₈ S ₃	7.49	0.59	6.7	32
24	V ₈ S ₄	6.70	0.61	3.3	73
25	V ₈ S ₅	6.77	0.75	2.9	39
26	V ₉ S ₁	7.45	0.79	3.0	48
27	V ₉ S ₅	7.01	0.56	3.1	40
28	V ₁₀ S ₂	7.78	0.64	5.8	35
29	V ₁₀ S ₄	7.86	0.54	4.6	20
30	V ₁₁ S ₂	7.52	0.87	5.3	43
31	V ₁₂ S ₃	7.72	0.86	7.6	57
32	V ₁₃ S ₂	7.60	0.63	3.7	32
33	V ₁₃ S ₃	7.58	0.73	6.9	58
34	V ₁₃ S ₄	7.51	0.68	4.6	70
35	V ₁₄ S ₁	7.12	0.27	4.8	35
36	V ₁₄ S ₂	6.95	0.54	4.6	41
37	V ₁₄ S ₃	7.58	0.33	4.1	78
38	V ₁₅ S ₁	7.52	0.76	6.5	55
39	V ₁₅ S ₂	7.48	0.63	3.1	70
40	V ₁₅ S ₃	7.62	0.35	5.1	57
41	V ₁₅ S ₄	7.64	0.58	4.4	60
42	V ₁₅ S ₅	7.51	0.39	5.1	66
43	V ₁₆ S ₂	7.57	0.77	5.4	72

44	V ₁₆ S ₄	7.58	0.66	3.7	70
45	V ₁₆ S ₅	7.61	0.72	1.5	35
46	V ₁₇ S ₃	7.63	0.83	3.3	65
47	V ₁₇ S ₅	7.55	0.49	4.2	75
48	V ₁₈ S ₂	8.14	0.31	2.3	40
49	V ₁₈ S ₃	7.72	0.41	5.2	35
50	V ₁₉ S ₁	7.60	0.83	4.8	63
51	V ₁₉ S ₂	7.82	0.79	5.1	59
52	V ₁₉ S ₄	7.60	0.45	4.0	57
53	V ₁₉ S ₅	7.94	0.47	4.5	82
54	V ₂₀ S ₁	7.62	0.69	3.8	49
55	V ₂₀ S ₄	8.02	0.34	4.5	57
56	V ₂₀ S ₅	8.09	0.83	6.2	67
Range		6.54-8.14	0.27-1.59	1.5 - 8.7	20-82
Mean		7.54	0.65	4.52	55.13
S.E.		0.04	0.03	0.18	2.07
C.V. (%)		4.16	37.07	30.69	28.13

Physico-chemical properties of Entisols in Renapur tahsil

Physico-chemical properties of Entisol soils are presented in table 2. From this data, it was revealed that the lowest soil pH value (6.65) was recorded in Khanapur village soil sample (V₁₀S₃) and the highest pH (8.11) was recorded from Wala village sample (V₁₈S₅). In this soil, pH was ranged from 4.36 to 8.11 with a mean value 7.51. The data presented in table 4 and depicted in fig 1 revealed that, out of 26 soil samples from Entisols, 4 samples (16%) were neutral and 22 samples (84%) were moderately alkaline in reaction.

The lowest EC (0.35 dSm⁻¹) was recorded in soil sample (V₁₈S₅) from Wala village, while the highest EC (1.25 dSm⁻¹) was observed from Sangvi village (V₆S₄). The range of EC was ranged from 0.35 to 1.25 dSm⁻¹ with an average value of 0.69 dSm⁻¹. The data presented in table 4 and depicted in fig 2 revealed that, out of 26 soil samples from Entisol, 23 samples (88%) were no deleterious effect on crop and 3 samples (12%) were critical for germination.

Organic carbon content was lowest (0.40 g kg⁻¹) in Arajkheda

village (V₅S₃) and the highest organic carbon content (7.80 g kg⁻¹) observed from Khanapur village (V₁₀S₁). The organic carbon content was ranged from 0.40 to 7.80 g kg⁻¹ with a mean value of 4.81 g kg⁻¹. The data presented in table 5 and depicted in fig 3 shows that out of 26 soil samples, 1 sample (4%) were very low, 8 samples (31%) were low, 11 samples (42%) were moderate and remaining 6 samples (23%) were moderately high content of organic carbon.

Lowest value of calcium carbonate (27 g kg⁻¹) was observed from Faradpur village (V₁₉S₃) and the highest value of calcium carbonate (119 g kg⁻¹) was recorded in soil sample (V₁₈S₅) from Wala village. The calcium carbonate content of Entisol soils ranged from 27 to 125 g kg⁻¹ with an average value of 81.39 g kg⁻¹. The data presented in table 5 and depicted in fig 4 revealed that, out of 26 soil samples from Entisol, 2 samples (8%) were Non calcareous (Fit for all crops), 20 samples (77%) were calcareous (Fit for all crops except citrus) and 4 samples (15%) were highly calcareous (Unfit for all crops) in nature.

Table 2: Physico-chemical properties of Entisol soils in Renapur tahsil

Sr. No.	Sample No.	pH	EC (dSm ⁻¹)	OC (g kg ⁻¹)	CaCO ₃ (g kg ⁻¹)
1	V ₅ S ₃	7.68	0.56	0.4	69
2	V ₅ S ₅	8.00	1.09	6.1	117
3	V ₆ S ₃	7.56	1.00	7.6	77
4	V ₆ S ₄	7.26	1.25	4.2	69
5	V ₇ S ₃	7.77	0.81	4.9	78
6	V ₉ S ₃	7.59	0.39	3.7	64
7	V ₁₀ S ₁	7.66	0.89	7.8	89
8	V ₁₀ S ₃	6.65	0.69	2.4	93
9	V ₁₀ S ₅	7.68	0.79	4.2	117
10	V ₁₁ S ₁	7.46	0.96	7.8	45
11	V ₁₁ S ₃	7.80	0.64	4.9	65
12	V ₁₁ S ₄	7.64	0.41	7.2	84
13	V ₁₂ S ₂	7.68	0.47	2.8	77
14	V ₁₂ S ₄	7.75	0.49	4.7	89
15	V ₁₂ S ₅	7.75	0.59	3.8	58
16	V ₁₃ S ₁	6.70	0.43	2.9	92
17	V ₁₃ S ₅	7.03	0.36	3.9	54
18	V ₁₄ S ₅	7.72	0.87	3.2	99
19	V ₁₆ S ₁	7.65	0.81	3.4	112
20	V ₁₆ S ₃	7.52	0.64	5.8	83
21	V ₁₇ S ₁	7.59	0.68	4.5	70
22	V ₁₇ S ₂	7.78	0.66	6.0	98
23	V ₁₈ S ₁	7.78	0.86	5.9	89
24	V ₁₈ S ₅	8.11	0.35	6.9	119
25	V ₁₉ S ₃	7.95	0.49	5.9	27
26	V ₂₀ S ₃	7.98	0.79	4.2	87

Range	6.65-8.11	0.35-1.25	0.4-7.8	27-119
Mean	7.51	0.69	4.81	81.39
S.E.	0.13	0.04	0.36	4.48
C.V. (%)	9.40	34.41	38.10	28.07

Physico-chemical properties of Vertisol soils in Renapur tahsil

The data of physico-chemical characteristics of Vertisols are presented in table 3. The lowest pH (7.00) was recorded in Harwadi soil sample (V_1S_3) while the highest pH (8.2) observed from Itti village ($V_{12}S_1$). The pH of the soils were ranged from 7.0 to 8.2 with an average value of 7.60. The data presented in table 4 and depicted in fig 1 noticed that out of 18 soil samples from Vertisols, 6 samples (34%) were neutral and 12 samples (66%) were moderately alkaline in reaction.

The highest EC (0.88 dSm^{-1}) was observed in soil samples of Nagapur ($V_{11}S_5$) and Harwadi (V_1S_3), while the lowest EC (0.28 dSm^{-1}) recorded in Harwadi village soil sample (V_1S_4). The EC value of soil sample were varied from 0.28 to 0.88 dSm^{-1} with a mean value 0.51 dSm^{-1} . The data presented in table 4 and depicted in fig 2 revealed that out of 18 soil samples from Vertisols, all 18 samples (100%) has no deleterious effect on crop.

The highest organic carbon content (7.9 g kg^{-1}) was observed in Sarola village ($V_{20}S_2$) and lowest organic carbon (2.2 g kg^{-1}) was observed in Shelu (V_2S_5) and Harwadi (V_1S_3) village. The range of organic carbon in these soil sample were 2.2 to 7.9 g kg^{-1} with a mean value 4.08 g kg^{-1} . The data presented in table 5 depicted in fig 3 revealed that, out of 18 samples, 10 samples (55%) were low, 7 samples (39%) moderate and remaining 1 sample (6%) was moderately high content of organic carbon.

The lowest value of calcium carbonate content (35 g kg^{-1}) was recorded from Harwadi village sample (V_1S_3) and the highest value calcium carbonate (110 g kg^{-1}) observed in Itti village sample ($V_{12}S_1$). Calcium carbonate content were varied from 35 to 110 g kg^{-1} with an average 63.65 g kg^{-1} . The data presented in table 5 and depicted in fig 4 noticed that out of 18 soil samples, 5 sample (28%) were non-calcareous (Fit for all crops), 12 samples (67%) were calcareous (Fit for all crops except citrus) and remaining 1 sample (5%) highly calcareous (Unfit for all crops) in nature.

The pH of Inceptisol, Entisol and Vertisol soils (Table 1, 2, 3) ranged from 6.54 to 8.14, 6.65 to 8.11 and 7 to 8.2 with a mean value 7.54, 7.51 and 7.60, respectively. The data presented in table 4 and depicted in fig 1 revealed that the 25, 16 and 34 per cent soil samples in Inceptisol, Entisol and Vertisol were neutral in reaction, respectively and 75, 84 and 66 per cent soil samples in Inceptisol, Entisol and Vertisol were moderately alkaline in reaction. These values of pH indicates that most of the soils under study were neutral to moderately alkaline in reaction. The alkaline reaction of soil is probably due to the presence of sufficient free lime content and basaltic alluvium parent material rich in aluminosilicate and alkaline earth from which these are derived. (Challa, 1995) [3]. The neutral nature of soil due to presence of bases and calcium/magnesium carbonate in the soils Borkar *et al.* (2018) [2]. The alkaline nature of soil is due to presence of free lime in the soils. Slightly alkaline nature of pH due to calcareous nature of these soils Magare *et al.* (2019) [8]. These results are also in close agreement with those reported by Patil *et al.* (2019) [10] noted pH range 4.5 to 8.1 from the soils of Washi tahsil of Osmanabad district.

From the data of table no., 1, 2, 3 it was observed that the EC

of the soils in Renapur tahsil ranged from 0.27 to 1.59 dSm^{-1} , 0.35 to 1.25 dSm^{-1} and 0.28 to 0.88 dSm^{-1} in Inceptisol, Entisol and Vertisol, respectively, with an average value of 0.65, 0.69 and 0.51 dSm^{-1} , respectively. The data presented in table 4 and depicted in fig 2 noticed that, 94 per cent, 88 per cent and 100 per cent soil samples of Inceptisol, Entisol and Vertisol were no deleterious effect on crop and 6 per cent, 12 per cent soil samples of Inceptisol, Entisol were critical for germination. The low content of EC was due to leaching of salt from the surface layer of soil. The values of EC obtained in analysis were found in desirable range as proposed by Richard and Cambell (1948) [12], when EC exceed 4 dSm^{-1} , the salt present become harmful to the crop growth. Ajaonkar and Patil (2017) [1] reported that the EC were ranged from 0.20 to 1.70 dSm^{-1} from the soils of Aurangabad district. These values of EC are safe for crop growth.

The organic carbon content in Inceptisol, Entisol and Vertisol soils (Table 1, 2, 3) ranged from 1.5 to 1.59, 0.4 to 7.8 and 2.2 to 7.9 g kg^{-1} with average value 4.52, 4.81 and 4.08 g kg^{-1} , respectively. Soils of these tahsil were low to medium in organic carbon content. The data presented in table 5 and depicted in fig 3 reported that, 41 per cent, 31 per cent and 55 per cent samples of Inceptisol, Entisol and Vertisol soils were low organic carbon content, respectively, while 12 per cent, 23 per cent and 6 per cent soil samples of Inceptisol, Entisol and Vertisol were moderately high in organic carbon content. The variation in organic carbon content in the soil due to high temperature which is responsible for hasten the rate of oxidation and very little addition of organic matter and crop residues in the soil. low to medium status of organic carbon was found in surveyed areas soil samples. The content of organic carbon in soils depends on the range of precipitation within experiment area, pattern of rainfall in the area. Organic carbon is also attributed to the variation in decomposition rate. Kashiwar *et al.* (2019) [7] noticed that the organic carbon content ranged from 2.7 to 8.6 g kg^{-1} from the soils of Sakoli tahsil of Bhandara district. Naiknaware (2018) [9] reported that the organic carbon ranged from 4.8 to 8.6 g kg^{-1} from the soils of Osmanabad district of Maharashtra.

The calcium carbonate content in Inceptisol, Entisol and Vertisol soils (Table 1, 2, 3) varied from 20 to 82, 27 to 119 and 35 to 110 g kg^{-1} with a mean value 55.13, 81.39 and 63.65 g kg^{-1} , respectively. The data presented in table 5 and depicted in fig 4 revealed that, 41 per cent, 8 per cent and 28 per cent soil samples of Inceptisol, Entisol and Vertisol were non-calcareous (Fit for all crops) in nature along with 59 percent, 77 per cent and 67 per cent soil samples of Inceptisol, Entisol and Vertisol were calcareous (Fit for all crops except citrus) in nature and 15 per cent and 5 per cent soil samples of Entisol and Vertisol were highly calcareous (Unfit for all crops) in nature. Data on calcium carbonate content of soils revealed that the soils were non-calcareous to calcareous and highly calcareous in nature. This might be due to presence of calcium carbonate in powdery form and hyper thermic temperature. Relative more accumulation of calcium carbonate in Vertisol soils and associated black soils, may be partly associated with their origin with rich in alkali earth and due to calcification process in this region. Waghmare *et al.* (2008) [13] reported that the CaCO_3 content in AUSA tahsil of

Latur district ranged from 8.80 to 125 g kg⁻¹. Ganorkar and Chinchmalatpure (2013) [4] studied soils of Rajura Bazar of Amravati district and noticed that the calcium carbonate

content were ranged from to 52.5 to 72.5 g kg⁻¹. Jibhakate *et al.* (2009) [6] reported that calcium carbonate content ranged from 38.4 to 96.9 g kg⁻¹ from Ketol tahsil of Nagpur district.

Table 3: Physico-chemical properties of Vertisol soils in Renapur tahsil

Sr. No.	Sample No.	pH	EC (dSm ⁻¹)	OC (g kg ⁻¹)	CaCO ₃ (g kg ⁻¹)
1	V ₁ S ₁	7.57	0.36	2.8	44
2	V ₁ S ₂	7.22	0.52	3.7	70
3	V ₁ S ₃	7.00	0.88	5.4	35
4	V ₁ S ₄	7.31	0.28	2.2	70
5	V ₁ S ₅	7.60	0.62	3.9	67
6	V ₂ S ₁	7.74	0.68	5.6	45
7	V ₂ S ₂	7.63	0.37	3.1	75
8	V ₂ S ₃	7.60	0.36	2.3	47
9	V ₂ S ₄	7.24	0.38	3.0	78
10	V ₂ S ₅	7.50	0.33	2.2	80
11	V ₉ S ₂	7.66	0.66	6.0	67
12	V ₉ S ₄	7.47	0.55	2.9	59
13	V ₁₁ S ₅	7.77	0.88	3.1	67
14	V ₁₂ S ₁	8.20	0.38	4.6	110
15	V ₁₄ S ₄	7.67	0.55	4.2	40
16	V ₁₇ S ₄	7.85	0.60	4.9	50
17	V ₁₈ S ₄	8.09	0.47	5.8	68
18	V ₂₀ S ₂	7.69	0.38	7.9	56
Range		7-8.2	0.28-0.88	2.2-7.9	35-110
Mean		7.60	0.51	4.08	63.65
S.E.		0.06	0.04	0.37	4.24
C.V. (%)		3.87	35.02	38.79	28.32

Table 4: Order wise categorization of soils from Renapur tahsil on the basis of pH and EC

pH	Soil orders	Categories	Strongly acidic (< 5.5)	Moderately acidic (5.5-6.0)	Slightly acidic (6.0-6.5)	Neutral (6.5-7.5)	Mod. alkaline (7.5-8.5)	Strong. alkaline (> 8.5)
	pH	Inceptisol	No. of sample	-	-	-	14	42
Per cent			-	-	-	25	75	-
Entisol		No. of sample	-	-	-	4	22	-
		Per cent	-	-	-	16	84	-
Vertisol		No. of sample	-	-	-	6	12	-
		Per cent	-	-	-	34	66	-
EC (dSm ⁻¹)	Soil orders	Categories	No deleterious effect on crop (< 1.0)		Critical for germination (1.0-2.0)	Critical for salt sensitive crop (2.0-3.0)		Injuries to most crops (> 3.0)
	Inceptisol	No. of sample	53		3	-		-
		Per cent	94		6	-		-
	Entisol	No. of sample	23		3	-		-
		Per cent	88		12	-		-
	Vertisol	No. of sample	18		-	-		-
		Per cent	100		-	-		-

Table 5: Order wise categorization of soil of Renapur tahsil on the basis of Organic carbon and CaCO₃ content

Organic carbon (g kg ⁻¹)	Soil orders	Categories	Very low (< 2)	Low (2.1-4.0)	Moderate (4.1-6.0)	Mod. High (6.1-8.0)	High (8.1-10)	Very high (> 10)
	Organic carbon (g kg ⁻¹)	Inceptisol	No. of sample	1	23	23	7	2
Per cent			2	41	41	12	4	-
Entisol		No. of sample	1	8	11	6	-	-
		Per cent	4	31	42	23	-	-
Vertisol		No. of sample	-	10	7	1	-	-
		Per cent	-	55	39	6	-	-
Calcium carbonate (g kg ⁻¹)	Soil orders	Categories	Non. Calcareous (< 50) (Fit for all crops)		Calcareous (50-100) (Fit for all crops except citrus)		Highly Calcareous (> 100) (Unfit for all crops)	
	Inceptisol	No. of sample	23		33		-	
		Per cent	41		59		-	
	Entisol	No. of sample	2		20		4	
		Per cent	8		77		15	
	Vertisol	No. of sample	5		12		1	
		Per cent	28		67		5	





Fig 1: Categorisation of soil pH in Renapur tahsil.






Fig 2: Categorization of EC in Renapur tahsil.




Fig 3: Categorization of organic carbon content in Renapur tahsil.




Fig 4: Categorisation of calcium carbonate content in soils of Renapur tahsil.

Conclusion

On the basis of soil characteristics, the soils from Renapur tahsil were grouped into Inceptisol, Entisol and Vertisol. In Renapur tahsil 56, 26 and 18 per cent soil samples were distributed into Inceptisol, Entisol and Vertisol, respectively. Most of the soils of Renapur tahsil was found neutral to moderately alkaline in reaction. EC of the soil was safe (no deleterious effect on crop). The organic carbon status of Renapur tahsil soil was low to moderate. Calcium carbonate content in these soils were non-calcareous (Fit for all crops) to calcareous (Fit for all crops except citrus) in nature.

References

1. Ajgaonkar SS, Patil SS. Soil health of soils in Aurangabad district (Maharashtra), India. *An Asian Journal of Soil Science*. 2017;12(1):121-127.
2. Borkar VS, Gokhale NB, Dhopakar RV, Khobragade NH, More SS, Kasture MC. Distribution of nutrient in different soil type in Kokan region of Maharashtra. *International Journal of Chemical Studies*. 2018;6(1):275-279.
3. Challa O, Vadivelu S, Sehgal JT. Soils of Maharashtra for optimizing land use. NBSS Pub 54 (soils of India series). NBSS and land use planning Nagpur, India, 1995, 112.
4. Ganorkar RP, Chinchmalapur PG. Physicochemical assessment of soil in Rajura bazar in Amravati district of Maharashtra (India). *International Journal of Chemical, Environmental and Pharmaceutical Research*. 2013;4(3):46-49.
5. Jackson ML. *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd. New Delhi, 1978.
6. Jibhakte SB, Bhende SN, Kharche VK, Selvalakshmi V. Physico-chemical status of soil of Katol tahsil in Nagpur district. *Journal of Soils and Crops*. 2009;19(1):122-128.
7. Kashiwar SR, Kundu MC, Dongarwar US. Assessment and mapping of soil nutrient status of Sakoli tahsil of Bhandara district of Maharashtra using GIS techniques. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(5):1900-1905.
8. Magare PN, Tedia K, Katkar RN, Lakhera ML, Shrivastava GK, Bajpai RK, *et al*. Physico-chemical

- characteristics and status of available micro and secondary nutrient in soils of agro-ecological-sub-region Latur district of Maharashtra. *Journal of Pharmacognosy and Phytochemistry*. 2019;8(5):1129-1134.
9. Naiknaware VV. Physico-chemical analysis of soil sample in Osmanabad district, Maharashtra. *National Conference on New Horizon in Material Science*. 2018;4(3):2394-4099.
 10. Patil PB, Jatkar PR, Adsul PB. Status of available nutrient of soils under vertisols from Washi tahsil of Osmanabad district of Maharashtra, India. *International Journal of Multidisciplinary Research and Development*. 2019;6(7):92-99.
 11. Piper CS. *Soil and plant analysis*. Hans Publication, Bombay, 1966.
 12. Richard H, Cambell P. In diagnosis and improvement of saline and alkali soils. *Hand book*, US. Dept. Agri, 1948, 60.
 13. Waghmare MS, Indulkar BS, Mali CV, Takankhar VG, Bavalgave VG. Chemical properties and micronutrient status of some soils of AUSA tahsil of Latur. *An Asian Journal of Soil Science*. 2008;3(2):236-241.